arXiv:0811.1047v1 [math.AG] 6 Nov 2008

INTRODUCTION TO THE MINIMAL MODEL PROGRAM AND THE EXISTENCE OF FLIPS

ALESSIO CORTI, ANNE-SOPHIE KALOGHIROS, AND VLADIMIR LAZIĆ

Abstract. The first aim of this note is to give a concise, but complete and self-contained, presentation of the fundamental theorems of Mori theory—the nonvanishing, base point free, rationality and cone theorems—using modern methods of multiplier ideals, Nadel vanishing, and the subadjunction theorem of Kawamata. The second aim is to write up a complete, detailed proof of existence of flips in dimension n assuming the minimal model program with scaling in dimension $n-1$.

Contents

1. Introduction 1
2. Basic definitions and results 3
3. Non-Vanishing, Basepoint Free and Rationality theorems 9
4. Log terminal and log canonical models 16
5. Restricted algebras and adjoint algebras 22
6. Shokurov's saturation condition 33
7. Existence of flips 36
References 37

1. Introduction

Our first aim in this note is to give a concise, but complete and self-contained, presentation of the fundamental theorems of Mori theory—the nonvanishing, base point free, rationality and cone theorems—using modern methods of multiplier ideals, Nadel vanishing, and the subadjunction theorem of Kawamata. We also give the basic definitions of log terminal and log canonical singularities of pairs, and spell out the minimal model program. We hope that the text can be used as a fast introduction to the field for those who wish quickly to master the foundations and equip themselves to do research. The approach here is not, at heart, different from the traditional one, but it is more efficient and it allows one to focus attention on the basic issues without being distracted by the technicalities. It also serves as a demonstration of the power of multiplier ideals, which play a crucial role in the most recent advances in the field, and the subadjunction theorem, which is a beautiful and increasingly central result.

We briefly sketch the key point as it arises in the proof of the base point free theorem. Let X be a nonsingular projective variety over $\mathbb{C}$, and L a nef divisor on X. Assuming that $L - \varepsilon K$ is ample for all small $\varepsilon > 0$, we want to show that all large multiples nL are base point free. The first step is to construct an effective $\mathbb{Q}$-divisor $D \sim mL$ (for some m) such that the pair (X, D) is not klt. For large n, we write $nL = K + D + A$, where A is ample. If c is largest such that (X, cD) is log canonical, then we write

$$nL = K + cD + A'$$

where $A' = A + (1 - c)D$ is still ample. Now, if cD contains a *divisor* S with multiplicity 1, then we may assume by induction on the dimension—and working with the pair $(S, cD_{|S})$—that all large multiples $nL_{|S}$ are base point free; standard vanishing theorems then imply that $H^0(X, nL) \to H^0(S, nL_{|S})$ is surjective, and we get the result. In general, the pair (X, cD) has some non-klt centres but they may all have higher codimension. The traditional approach is to blow up until one of the centres becomes a divisor (in fact, one blows up indiscriminately to a log resolution of everything in sight). Instead, in these notes, we use multiplier ideals, Nadel Vanishing, and the subadjunction theorem of Kawamata, to work *directly on* X and lift sections from a minimal non-klt centre $W \subset X$ of the pair (X, cD).

Our second aim is to write up a complete, detailed proof of existence of flips in dimension n assuming the minimal model program with scaling in dimension $n - 1$. Our proof is closer in spirit to some of Shokurov's original arguments in [Sho03] than to the treatment of [HM05] and, especially, [HM08]. Our aim is to present as *robust* a proof as possible. In short, we proceed as follows. The starting point is a pl flipping contraction $f\colon (X, S + B) \to Z$:

- X is n-dimensional and the pair $(X, S+B)$ has plt singularities;
- f is a flipping contraction for $K + S + B$;
- S is non-empty and f-negative.

The aim is to construct the flip of f by showing that the canonical algebra

$$R = R(X, K + S + B) = \bigoplus_{n=0}^{\infty} H^0\big(X, n(K + S + B)\big)$$

is finitely generated. Shokurov's great insight was to suggest that we can do this by showing that the *restricted algebra*

$$R_S = \mathrm{Image}\big(R \to k(S)[T]\big)$$

is finitely generated. (R is an algebra of rational functions and we get R_S by restricting those rational functions to S.) We develop Shokurov's language of b-divisors to make sense of the statement that R_S is a b-divisorial algebra. We then show that R_S enjoys two key properties:

1. R_S is an *adjoint algebra*—this notion comes from a further major insight of Hacon and M$^{\mathrm{c}}$Kernan and the proof of this fact uses their important lifting lemma;
2. R_S is *saturated* in the sense of Shokurov.

Using these two properties, and the minimal model program with scaling in dimension $n-1$, we can then show that R_S is finitely generated. Our argument is robust in the sense that the two key properties—adjointness and saturation—are treated as separate issues and proved independently of each other.

This note is a cleaned-up version of AC's lectures at the Summer School in Grenoble *Geometry of complex projective varieties and the minimal model program* 18 June–06 July 2007. We thank the organisers of that very successful event for providing an ideal setting and a comprehensively positive atmosphere.

It is our pleasure to acknowledge the influence of Robert Lazarsfeld on the point of view endorsed in these notes. It was he who re-iterated Kawamata's suggestion that a more transparent proof of the fundamental theorems of Mori theory is possible based on multiplier ideals and the subadjunction theorem. We especially thank Alex Küronya for his careful reading of earlier versions of these notes and his useful comments. We also thank Christopher Hacon and Sándor Kovács.

We tried hard to chase down and remove mistakes from the text; please accept our apologies for those that must inevitably be still around.

2. Basic definitions and results

Notation 2.1. All varieties in this paper are proper, irreducible and normal over $\mathbb{C}$. We mostly work with projective varieties. We write $\sim$ for linear equivalence of Weil divisors and $\equiv$ for numerical equivalence of Cartier divisors. On a variety X, $\mathrm{WDiv}(X)$ denotes the group of Weil divisors, $\mathrm{Div}(X)$ the group of Cartier divisors and $\mathrm{Pic}(X) = \mathrm{Div}(X)/\sim$. Subscripts denote either the ring in which the coefficients of divisors are taken or that the equivalence is relative to a specified morphism. An ample $\mathbb{Q}$-divisor A is *general* if kA is a general member of the linear system $|kA|$ for some $k \gg 0$.

Definition 2.2.

1. A *log pair* (X, Δ) consists of a variety X and a divisor $\Delta \in \mathrm{WDiv}(X)_{\mathbb{R}}$ such that $K_X + \Delta$ is $\mathbb{R}$-Cartier.
2. A pair (X, Δ) is *log smooth* if X is nonsingular and $\operatorname{Supp} \Delta$ has simple normal crossings.
3. A *model* over X is a proper birational morphism $f\colon Y \to X$.
4. A *log resolution* of (X, Δ) is a model $f\colon Y \to X$ such that $(Y, f_*^{-1}\Delta + \operatorname{Exc} f)$ is log smooth.
5. A *boundary* is a divisor $\Delta = \sum d_i D_i \in \mathrm{WDiv}(X)_{\mathbb{R}}$ such that $0 \leq d_i \leq 1$ for all i.

Pairs first arose through the study of open varieties (Iitaka program). If U is quasi-projective, and if X is a compactification of U such that $\Delta = X \setminus U$ has simple normal crossings, then the ring

$$R(X, \Delta) = \bigoplus_{n \in \mathbb{N}} H^0(X, n(K_X + \Delta))$$

depends only on the open variety U [Iit82, Chapters 10 and 11].

A valuation $\nu\colon k(X) \to \mathbb{Z}$ is *geometric* if $\nu = \mathrm{mult}_E$, where $E \subset Y$ is a prime divisor in a model $Y \to X$. Divisors $E \subset Y \to X$ and $E' \subset Y' \to X$ define the same geometric valuation if and only if the induced birational map $Y \dashrightarrow Y'$ is an isomorphism at the generic points of E and E'. The *centre* of a geometric valuation ν on X associated to a divisor E on a model $f\colon Y \to X$ is denoted by $c_X\nu = f(E)$. A geometric valuation ν is *exceptional* when $\operatorname{codim}_X(c_X\nu) \geq 2$. We often identify a geometric valuation ν and the corresponding divisor E.

Definition 2.3. An integral *b-divisor* $\mathbf{D}$ on X is an element of the group

$$\mathbf{Div}(X) = \varprojlim \mathrm{WDiv}(Y),$$

where the limit is taken over all models $f\colon Y \to X$ with the induced homomorphisms $f_*\colon \mathrm{WDiv}(Y) \to \mathrm{WDiv}(X)$. Thus $\mathbf{D}$ is a collection of divisors $\mathbf{D}_Y \in \mathrm{WDiv}(Y)$ compatible with push-forwards. Each $\mathbf{D}_Y$ is the *trace* of $\mathbf{D}$ on Y.

For every model $f\colon Y \to X$ the induced map $f_*\colon \mathbf{Div}(Y) \to \mathbf{Div}(X)$ is an isomorphism, so b-divisors on X can be identified with b-divisors on any model over X. For every open subset $U \subset X$ we naturally define the restriction $\mathbf{D}_{|U}$ of a b-divisor $\mathbf{D}$ on X.

Definition 2.4. The *b-divisor of a nonzero rational function* φ is

$$\mathbf{div}_X\,\varphi = \sum \nu_E(\varphi)E,$$

where E runs through geometric valuations with centre on X. Two b-divisors are *linearly equivalent* if they differ by the b-divisor of a nonzero rational function.

The associated *b-divisorial sheaf* $\mathcal{O}_X(\mathbf{D})$ is defined by

$$\Gamma(U, \mathcal{O}_X(\mathbf{D})) = \{\varphi \in k(X) : (\mathbf{div}_X\,\varphi + \mathbf{D})_{|U} \geq 0\}.$$

Definition 2.5. The *proper transform* b-divisor $\widehat{D}$ of an $\mathbb{R}$-divisor D has trace $\widehat{D}_Y = f_*^{-1}D$ on every model $f\colon Y \to X$.

The *Cartier closure* of an $\mathbb{R}$-Cartier divisor D on X is the b-divisor $\overline{D}$ with trace $\overline{D}_Y = f^*D$ on every model $f\colon Y \to X$.

A b-divisor $\mathbf{D}$ *descends* to a model $Y \to X$ if $\mathbf{D} = \overline{\mathbf{D}_Y}$; we then say that $\mathbf{D}$ is a *Cartier b-divisor*.

A b-divisor $\mathbf{D}$ on X is *good* on a model $Y \to X$ if $\mathbf{D} \geq \overline{\mathbf{D}_Y}$; in particular the sheaf $\mathcal{O}(\mathbf{D})$ is coherent.

A b-divisor $\mathbf{M}$ on X is *mobile* if it descends to a model $Y \to X$ where $\mathbf{M}_Y$ is basepoint free.

Note that if a mobile b-divisor $\mathbf{M}$ descends to a model $W \to X$, then $\mathbf{M}_W$ is free and $H^0(X, \mathbf{M}) = H^0(W, \mathbf{M}_W)$.

Lemma 2.6. *Let* $\mathbf{N}$ *be a good b-divisor on* X*. There is a mobile b-divisor* $\mathbf{M}$ *such that* $\mathbf{M}_Y = \operatorname{Mob}\mathbf{N}_Y$ *for every model* $f\colon Y \to X$ *and* $H^0(X, \mathbf{N}) = H^0(Y, \mathbf{N}_Y)$.

Proof. Since $\mathbf{N}_Y = f^*\mathbf{N}_X + E$ for some effective and exceptional divisor E, we have $f_*\operatorname{Mob}\mathbf{N}_Y = f_*\operatorname{Mob} f^*\mathbf{N}_X = \operatorname{Mob}\mathbf{N}_X$. ☐

Cartier restriction. Let $\mathbf{D}$ be a Cartier b-divisor on X and let S be a normal prime divisor in X such that $S \not\subset \operatorname{Supp}\mathbf{D}_X$. Let $f\colon Y \to X$ be a log resolution of (X, S) such that $\mathbf{D}$ descends to Y. Define the *restriction* of $\mathbf{D}$ to S as

$$\mathbf{D}_{|S} := \overline{\mathbf{D}_{Y|\widehat{S}_Y}}.$$

This is a b-divisor on S via $(f_{|\widehat{S}_Y})_*$ that does not depend on the choice of log resolution. By definition, $\mathbf{D}_{|S}$ is a Cartier b-divisor that satisfies $(\mathbf{D}_1 + \mathbf{D}_2)_{|S} = \mathbf{D}_{1|S} + \mathbf{D}_{2|S}$, and $\mathbf{D}_{1|S} \geq \mathbf{D}_{2|S}$ if $\mathbf{D}_1 \geq \mathbf{D}_2$.

Definition 2.7.

1. The *canonical* b-divisor $\mathbf{K}_X$ on X has trace $(\mathbf{K}_X)_Y = K_Y$ on every model $Y \to X$.

2. The *discrepancy* $\mathbf{A}(X,\Delta)$ of the pair (X,Δ) is
$$\mathbf{A}(X,\Delta)=\mathbf{K}_X-\overline{K_X+\Delta}.$$
3. For a geometric valuation E on X, the *discrepancy of* E with respect to (X,Δ) is
$$a(E,X,\Delta)=\operatorname{mult}_E\mathbf{A}(X,\Delta).$$

Definition 2.8. Let (X,Δ) be a pair and let D be an $\mathbb{R}$-Cartier divisor on X. The *multiplier ideal sheaf* associated to (X,Δ) and D is
$$\mathcal{J}((X,\Delta);D)=\mathcal{O}_X(\lceil\mathbf{A}(X,\Delta+D)\rceil).$$

Remark 2.9. We obviously have $\mathcal{J}((X,\Delta);D)=\mathcal{J}((X,\Delta+D);0)$. When no confusion is likely we write $\mathcal{J}(\Delta)=\mathcal{J}((X,\Delta);0)$.

Theorem 2.10 (Local vanishing)**.** *Let $f\colon Y\to X$ be a log resolution of a pair (X,Δ). Then*
$$R^if_*\mathcal{O}_Y(\lceil\mathbf{A}(X,\Delta)_Y\rceil)=0$$
for all $i>0$.

Proof. If X is projective, consider an ample divisor A such that for $m\gg0$, the divisors $mA-\Delta$ are ample, the sheaves $R^if_*\mathcal{O}_Y(K_Y-\lfloor f^*\Delta\rfloor)\otimes\mathcal{O}_X(mA)$ are globally generated (when non-zero), and
$$H^j(X,R^if_*\mathcal{O}_Y(K_Y-\lfloor f^*\Delta\rfloor)\otimes\mathcal{O}_X(mA))=(0)$$
for all $j>0$ and $i\geq0$. The Leray spectral sequence and Kawamata-Viehweg vanishing yield
$$\begin{aligned}H^0(X,R^if_*&\mathcal{O}_Y(K_Y-\lfloor f^*\Delta\rfloor)\otimes\mathcal{O}_X(mA))\\&=H^i(Y,\mathcal{O}_Y(K_Y-\lfloor f^*\Delta\rfloor+f^*(mA)))=(0).\end{aligned}$$
In particular, the sheaves $R^if_*\mathcal{O}_Y(K_Y-\lfloor f^*\Delta\rfloor)\otimes\mathcal{O}_X(mA)$ are zero because they are globally generated, and thus
$$R^if_*\mathcal{O}_Y(\lceil\mathbf{A}(X,\Delta)_Y\rceil)=R^if_*\mathcal{O}_Y(K_Y-\lfloor f^*\Delta\rfloor)\otimes\mathcal{O}_X(-K_X)=0$$
for $i>0$.

For the non projective case, see [Laz04, Theorem 9.4.1]. □

Theorem 2.11 (Nadel vanishing)**.** *Let L be a Cartier divisor on X such that $L-(K_X+\Delta)$ is nef and big. Then*
$$H^i\big(X,\mathcal{O}_X(L)\otimes\mathcal{J}(\Delta)\big)=(0)$$
for all $i>0$.

Proof. Let $f\colon Y \to X$ be a log resolution of (X,Δ). The Local vanishing theorem states that

$$R^i f_* \mathcal{O}_Y(\lceil \mathbf{A}(X,\Delta)_Y \rceil + f^*L) = 0$$

for $i > 0$; the Leray spectral sequence and Kawamata-Viehweg vanishing imply

$$H^i(X, \mathcal{O}_X(L) \otimes \mathcal{J}(\Delta))) = H^i(Y, \lceil \mathbf{A}(X,\Delta)_Y \rceil + f^*L) = (0)$$

for $i > 0$. $\square$

The following result is a characterisation of *big* divisors.

Definition-Lemma 2.12 (Kodaira's Lemma)**.** Let D be a Cartier divisor on a projective variety X of dimension n. The divisor D is *big* if one of the following equivalent conditions holds:

1. There exists $C > 0$ such that $h^0(X, kD) > Ck^n$ for $k \gg 1$.
2. For any ample divisor A on X, there exists $m \in \mathbb{N}$ and an effective divisor E on X such that $mD \sim A + E$.

If D is nef, D is big if and only if one of the following conditions holds:

1. $D^n > 0$.
2. There is an effective divisor E and ample $\mathbb{Q}$-divisors A_k such that $D \equiv A_k + (1/k)E$ for $k \gg 1$.

We recall some definitions of singularities of pairs.

Definition 2.13.

1. A log pair (X,Δ) is *Kawamata log terminal (klt)* if $\lceil \mathbf{A}(X,\Delta) \rceil \geq 0$, or equivalently if $\mathcal{J}(\Delta) = \mathcal{O}_X$.
2. A log pair (X,Δ) is *purely log terminal (plt)* if $a(E,X,\Delta) > -1$ for every exceptional geometric valuation E on X; in particular $\lceil \Delta \rceil$ is reduced and it can be proved that the connected components of $\lceil \Delta \rceil$ are normal.
3. A log pair (X,Δ) is *log canonical (lc)* if $a(E,X,\Delta) \geq -1$ for every geometric valuation E on X.
4. A log pair (X,Δ) is *divisorially log terminal (dlt)* if there is a log resolution $Y \to X$ such that $\lceil \mathbf{A}(X,\Delta)_Y + \widehat{\Delta}_Y \rceil \geq 0$.

Definition-Lemma 2.14. Let (X,Δ) be a lc pair.

1. The *non-klt locus* of (X,Δ) is
$$\operatorname{nklt}(X,\Delta) = \operatorname{Supp}\big(\mathcal{O}_X/\mathcal{J}(\Delta)\big).$$
2. A *non-klt centre* is the centre W of a geometric valuation such that $W \subset \operatorname{nklt}(X,\Delta)$; therefore $\mathcal{J}(\Delta) \subset \mathcal{I}_W$.

3. A non-klt centre W is *isolated* if for any geometric valuation E on X such that $a(E,X,\Delta)=-1$, $c_XE=W$. A non-klt centre W is *exceptional* if it is isolated and if there is a unique geometric valuation E on X with $a(E,X,\Delta)=-1$.
4. If W is an isolated non-klt centre, $\mathcal{J}(\Delta)=\mathcal{I}_W$.

Proof. Let W be an isolated non-klt centre and let $f\colon Y\to X$ be a log resolution. Then $\lceil \mathbf{A}(X,\Delta)_Y\rceil=-E+A$, where E and A are effective, each component of E maps onto W and A is f-exceptional. Thus $f_*\mathcal{O}_Y(A)=\mathcal{O}_X$ and since $R^1f_*\mathcal{O}_Y(-E+A)=0$ by Local vanishing, applying f_* to the sequence

$$0\to\mathcal{O}_Y(-E+A)\to\mathcal{O}_Y(A)\to\mathcal{O}_E(A_{|E})\to 0$$

yields

$$0\to\mathcal{J}(\Delta)\to\mathcal{O}_X\to f_*\mathcal{O}_E(A_{|E})\to 0.$$

Therefore $f_*\mathcal{O}_E(A_{|E})$ is a quotient of $\mathcal{O}_X$ which is an $\mathcal{O}_W$-sheaf since $f(E)=W$, so $f_*\mathcal{O}_E(A_{|E})=\mathcal{O}_W$ and $\mathcal{J}(\Delta)=\mathcal{I}_W$. ☐

Remark 2.15. Let $f\colon Y\to X$ be a log resolution of (X,Δ) and for $1\leq i\leq m$, let $E_i\subset Y$ be the geometric valuations on X with $a(E_i,X,\Delta)=-1$. The non-klt centres of (X,Δ) are precisely the images $f(\cap_{i\in N}E_i)$ for $N\subset\{1,\dots,m\}$. In particular, for any $x\in\operatorname{nklt}(X,\Delta)$, there is a well defined *minimal non-klt centre* through x.

Lemma 2.16 (Tie breaking, [Kol07, Proposition 8.7.1])**.** *Let (X,Δ) be a klt pair and D a $\mathbb{Q}$-Cartier divisor on X such that $(X,\Delta+D)$ is lc. Let W be a minimal non-klt centre of $(X,\Delta+D)$ and let A be an ample divisor. Then there are arbitrarily small $\varepsilon,\eta>0$ and a divisor $D'\sim_{\mathbb{Q}}A$ such that W is an exceptional non-klt centre of $(X,\Delta+(1-\varepsilon)D+\eta D')$.*

Proof. We show how to make the non-klt centre W isolated. Let $f\colon Y\to X$ be a log resolution of the pair $(X,\Delta+D)$. We write

$$K_Y=f^*(K_X+\Delta+D)+\sum a_iE_i\quad\text{and}\quad f^*D=\sum b_iE_i,$$

so that

$$K_Y=f^*(K_X+\Delta+(1-\varepsilon)D)+\sum(a_i+\varepsilon b_i)E_i.$$

Since $(X,\Delta+D)$ is lc and (X,Δ) is klt, $a_i\geq-1$, $a_i+b_i>-1$ and hence $a_i+\varepsilon b_i>-1$ for all $\varepsilon>0$.

Let Z be a Cartier divisor such that W is the only non-klt centre contained in $\operatorname{Supp}Z$ and let D_0 be a general member of the linear system $|pA-Z|$ for a large p. Consider a $\mathbb{Q}$-divisor $D'=\frac{1}{p}(Z+D_0)\sim_{\mathbb{Q}}A$; in particular W is the only non-klt centre contained in $\operatorname{Supp}D'$. Write $f^*D'=\sum d_iE_i$, where we may assume that f is a log resolution of $(X, \Delta + D + D')$. By construction, if $a_i = -1$ then $d_i > 0$ precisely when $f(E_i) = W$.

The pair $(X, \Delta + (1 - \varepsilon)D + \eta D')$ is lc if and only if

$$a_i + \varepsilon b_i - \eta d_i \geq -1$$

for all i. If $a_i > -1$, this holds for all sufficiently small $\varepsilon, \eta > 0$. If $a_i = -1$ and $f(E_i) \neq W$, then $a_i + \varepsilon b_i - \eta d_i > -1$ for all $\varepsilon > 0$. Fix ε and define

$$\eta = \min\{\varepsilon b_i / d_i : a_i = -1, d_i > 0\}.$$

By construction, the pair $(X, \Delta + (1 - \varepsilon)D + \eta D')$ is log canonical and W is an isolated non-klt centre.

The non-klt centre W can further be made exceptional as in [Kol07, Proposition 8.7.1]. $\square$

Theorem 2.17 (Kawamata's Subadjunction, [Kol07, Theorem 8.6.1])**.** *Let (X, Δ) be an lc pair. Assume that W is an exceptional lc centre of (X, Δ) and let A be an ample $\mathbb{R}$-divisor. Then W is normal and for every $\varepsilon > 0$,*

$$(K_X + \Delta + \varepsilon A)_{|W} \sim_{\mathbb{R}} K_W + \Delta_W$$

for some divisor Δ_W on W such that the pair (W, Δ_W) is klt.

Remark 2.18. The choice of the divisor Δ_W in its linear equivalence class is not canonical.

Theorem 2.19. *Let $(X, S + B)$ be a plt pair, where S is irreducible. Then*

$$(K_X + S + B)_{|S} \sim_{\mathbb{R}} K_S + B_S$$

for some divisor B_S on S such that the pair (S, B_S) is klt.

Remark 2.20. Our notation is not the same as the one used in [KM98], where B_S is called the *different* and denoted $\mathrm{Diff}(B)$.

Definition 2.21. Let (X, Δ) be a pair and let x be a point in X. The *log canonical threshold* of (X, Δ) at x is

$$\mathrm{lct}(\Delta; x) = \sup\{c \in \mathbb{R} : (X, c\Delta) \text{ is lc in the neighbourhood of } x\}.$$

3. Non-Vanishing, Basepoint Free and Rationality theorems

We present the Minimal Model Program (MMP) in a relative setting, that is, given a birational projective morphism $f\colon X \to Z$ we consider the MMP over Z. The motivation for this is clear when, for example, X has a fibration structure over Z.

Theorem 3.1 (Cone Theorem)**.** *Let (X, Δ) be a dlt pair over Z. Then there are $(K_X + \Delta)$-negative rational curves $C_i \subset X$ such that*

$$\overline{NE}(X/Z) = \overline{NE}(X/Z)_{K_X+\Delta\geq 0} + \sum R_i,$$

where the $(K_X + \Delta)$-negative extremal rays *R_i are spanned by the classes of C_i and are locally discrete in $\overline{NE}(X/Z)_{K_X+\Delta<0}$.*

Theorem 3.2 (Contraction Theorem)**.** *Let (X, Δ) be a dlt pair over Z. Let $R \subset \mathrm{NE}(X/Z)$ be a $(K_X + \Delta)$-negative extremal ray. There is a morphism $\varphi\colon X \to X'$ that is characterised by:*

1. *$\varphi_*\mathcal{O}_X = \mathcal{O}_{X'}$,*
2. *an effective curve is contracted by φ if and only if its class belongs to R.*

The morphism φ is the contraction of R.

Remark 3.3. These two theorems hold for lc pairs [Amb03].

Theorem 3.4 (Basepoint Free Theorem)**.** *Let (X, Δ) be a klt pair and let L be a nef Cartier divisor on X. Assume that there exists $p > 0$ such that $pL - (K_X + \Delta)$ is nef and big. Then the linear system $|nL|$ is basepoint free for all $n \gg 0$.*

Remark 3.5. The Zariski counterexample [Zar62] shows that the Basepoint Free theorem does not hold for dlt pairs when $pL - (K_X + \Delta)$ is assumed to be nef and big. It does hold however when $pL - (K_X + \Delta)$ is ample.

Corollary 3.6. *Let $\varphi\colon X \to Y$ be the contraction of an extremal ray R. The sequence*

$$0 \longrightarrow \mathrm{Pic}(Y) \xrightarrow{\varphi^*} \mathrm{Pic}(X) \longrightarrow \mathbb{Z}$$

is exact, where the last map in the sequence is multiplication by a fixed curve C whose class belongs to R. If $L \cdot C = 0$, then $|nL|$ is basepoint free.

Definition-Lemma 3.7. Let (X, Δ) be a $\mathbb{Q}$-factorial dlt pair, and let $\varphi\colon X \to Y$ be the contraction of a $(K_X + \Delta)$-negative extremal ray R.

1. If $\dim Y < \dim X$, then φ is a *Mori fibration*;
2. If $\dim Y = \dim X$ and $\mathrm{Exc}(\varphi)$ is an irreducible divisor, then φ is a *divisorial contraction*;
3. If $\dim Y = \dim X$ and $\operatorname{codim}_X \operatorname{Exc} \varphi \geq 2$, i.e. φ is a *small* map, then φ is a *flipping contraction.*

Proof. The only thing there is to prove is that in the second case, if $E \subset \operatorname{Exc}(\varphi)$ is a prime divisor, then $E = \operatorname{Exc}(\varphi)$. Let C be any curve that is contracted by f. As the class of C belongs to R, $E \cdot C < 0$ by Lemma 3.9, and $C \subset E$. □

Remark 3.8. If $\varphi\colon X \to Y$ is a small extremal contraction, Y is not $\mathbb{Q}$-factorial. In higher dimensions, a surgery operation in codimension 2 is necessary to proceed with the MMP. This operation is the *flip* of φ; it is defined in Section 4.

Lemma 3.9 (Negativity Lemma, [KM98, Lemma 3.38])**.** *Let $f\colon Y \to X$ be a birational morphism and let $E = \sum a_i E_i$ be an f-exceptional divisor on Y. Assume that*

$$E \equiv_f H + D,$$

where H is f-nef and D is an effective divisor that has no common components with E. Then all $a_i \leq 0$. If $x \in X$ is a point such that D or H is not numerically trivial on $f^{-1}(x)$, and if E_i is a divisor such that $f(E_i) = x$, then $a_i < 0$.

The following result is the first step in proving Theorem 3.4.

Theorem 3.10 (Non-Vanishing Theorem)**.** *Let (X, Δ) be a klt pair and let L be a nef Cartier divisor on X. Assume that there exists $p > 0$ such that $pL - (K_X + \Delta)$ is nef and big. Then the linear system $|nL|$ is non-empty for all $n \gg 0$.*

Proof. The proof is by induction on $d = \dim X$.

We may assume that $pL - (K_X + \Delta)$ is ample and that Δ is a $\mathbb{Q}$-divisor. Indeed, Definition-Lemma 2.12 shows that there is an effective divisor E such that for all $0 < \varepsilon \ll 1$, $pL - (K_X + \Delta) - \varepsilon E$ is ample, and $(X, \Delta + \varepsilon E)$ is klt since (X, Δ) is.

If L is numerically trivial, by Kawamata-Viehweg vanishing

$$h^0(X, nL) = \chi(\mathcal{O}_X(nL)) = \chi(\mathcal{O}_X) = h^0(\mathcal{O}_X) \neq 0$$

and the linear series $|nL|$ is non-empty for all $n \geq 0$.

Assume that L is not numerically trivial. First, we show that for $q \gg 1$,

$$qL \sim K_X + \Delta + A,$$

where A is an ample divisor and the pair $(X, \Delta + A)$ is not klt. Let C be a curve such that $L \cdot C \neq 0$. Since $pL - (K_X + \Delta)$ is ample, there is a positive integer m such that $(m(pL - (K_X + \Delta)))^{d-1}$ is represented by C plus an effective 1-cycle. Thus $L \cdot (pL - (K_X + \Delta))^{d-1} > 0$, and therefore the intersection number

$$\begin{aligned}(qL - (K_X + \Delta))^d &= ((q-p)L + pL - (K_X + \Delta))^d \\ &\geq (q-p)L \cdot (pL - (K_X + \Delta))^{d-1}\end{aligned}$$

tends to infinity with q. By Riemann-Roch and Kawamata-Viehweg vanishing, for $q \gg 0$,

$$h^0(X, n(qL - (K_X + \Delta))) = \frac{n^d}{d!}(qL - (K_X + \Delta))^d + O(d-1).$$

If $x \in X$ is a nonsingular point, a parameter count shows that there is a section $D_{q,n} \in |n(qL - (K_X + \Delta))|$ with $\operatorname{mult}_x D_{q,n} \geq (d+1)n$ for q, n sufficiently large.

Fix one such pair q, n and denote $A = \frac{1}{n} D_{q,n}$. By construction, $(X, \Delta + A)$ is not klt at x. Let

$$c = \min\{t \in \mathbb{R} : (X, \Delta + tA) \text{ is not klt}\} \leq 1$$

and consider a minimal non-klt centre W of $(X, \Delta + cA)$. By Lemma 2.16, we may assume that W is an exceptional non-klt centre of $(X, \Delta + cA)$.

For $0 < \varepsilon \ll 1$,

$$qL \sim_{\mathbb{Q}} K_X + \Delta + cA + \varepsilon A + (1 - c - \varepsilon)A,$$

and Theorem 2.17 shows that

$$qL_{|W} \sim_{\mathbb{Q}} K_W + \Delta_W + (1 - c - \varepsilon)A_{|W} \tag{1}$$

where (W, Δ_W) is a klt pair. As $qL_{|W}$ is nef and $qL_{|W} - (K_W + \Delta_W)$ is ample, by induction $|qL_{|W}| \neq \emptyset$. The non-klt centre W is exceptional, therefore by Definition-Lemma 2.14, $\mathcal{J}((X, \Delta); cA) = \mathcal{I}_W$. By Nadel vanishing,

$$H^i(X, \mathcal{I}_W(qL)) = (0)$$

for $i > 0$. The long exact sequence in cohomology associated to

$$0 \to \mathcal{I}_W(qL) \to \mathcal{O}_X(qL) \to \mathcal{O}_W(qL_{|W}) \to 0$$

yields a surjection $H^0(X, qL) \to H^0(W, qL_{|W})$, so the linear series $|qL|$ is non-empty. $\square$

Remark 3.11. When working with klt pairs, one can apply Kodaira's Lemma to replace nef and big divisors with ample ones. However, this no longer holds in the context of dlt pairs.

Proof of Theorem 3.4. By Theorem 3.10, the linear system $|nL|$ is non-empty for $n \gg 0$; we now show that $|nL|$ is in fact basepoint free for $n \gg 0$. As is explained in Remark 3.11, we may assume that $pL - (K_X + \Delta)$ is ample.

For a positive integer n, since $\operatorname{Bs}|n^uL| \subset \operatorname{Bs}|n^vL|$ when $u > v$, the Noetherian condition shows that the sequence $\operatorname{Bs}|n^uL|$ stabilises. Denote by B_n its limit. If $B_p = B_q = \emptyset$ for two relatively prime integers p and q, then $\operatorname{Bs}|p^{u_0}L| = \operatorname{Bs}|q^{v_0}L| = \emptyset$. Since every $n \gg 0$ can be written as a non-negative linear combination of p^{u_0} and q^{v_0}, $|nL|$ is basepoint free.

Assume that there is an integer m such that $B_m \neq \emptyset$. By taking a multiple of L, we may assume that $B_m = \operatorname{Bs}|mL| = \operatorname{Bs}|m^eL|$ for all $e \geq 1$. Let D be a general section of $|mL|$.

Define

$$c = \inf\{t \in \mathbb{R} : (X, \Delta + tD) \text{ is not klt}\} > 0,$$

so that the pair $(X, \Delta + cD)$ is strictly lc. Note that $c \leq 1$ because D is an integral divisor.

We choose an appropriate minimal non-klt centre of $(X, \Delta + cD)$ as follows. If there is a minimal non-klt centre W of codimension at least 2, then $W \subset \operatorname{Bs}|mL|$. If all minimal non-klt centres are of codimension 1, note that $c = 1$ and that every component of D is a minimal non-klt centre. Let W be a component of D that intersects $\operatorname{Bs}|mL|$.

By Lemma 2.16, we may assume that W is an exceptional non-klt centre of the pair $(X, \Delta + cD)$ and therefore $\mathcal{J}(X, \Delta + cD) = \mathcal{I}_W$ by Definition-Lemma 2.14.

Fix $q = m^v \geq p + m$. Since L is nef we have

$$qL \sim_{\mathbb{Q}} K_X + \Delta + D + A,$$

where A is an ample $\mathbb{Q}$-Cartier divisor. Then

$$qL \sim_{\mathbb{Q}} K_X + \Delta + cD + \varepsilon A + A_\varepsilon,$$

where $A_\varepsilon = (1-c)D + (1-\varepsilon)A$ is ample for ε sufficiently small. Theorem 2.17 then gives

$$qL_{|W} \sim_{\mathbb{Q}} K_W + \Delta_W + A_{\varepsilon|W}.$$

As in the proof of Theorem 3.10 the map

$$H^0(X, qL) \to H^0(W, qL_{|W})$$

is surjective, and therefore $\operatorname{Bs}|qL| \cap W = \operatorname{Bs}|qL_{|W}|$. But by induction on the dimension $\operatorname{Bs}|qL_{|W}| = \emptyset$ and that is a contradiction.

□

Remark 3.12 (Effective basepoint free theorem [Kol93])**.** Under the hypotheses of Theorem 3.4, one can show that there is a positive integer m that depends only on $\dim X$ and p such that $|nL|$ is basepoint free for $n \geq m$.

The following lemma will be used in the proof of the Rationality theorem.

Lemma 3.13 ([KM98])**.** *Let $P(x,y)$ be a non-trivial polynomial of degree at most n. Fix a positive integer a and a positive real number ε. Assume that P vanishes for all sufficiently large $x,y \in \mathbb{N}$ such that $0 < ay - rx < \varepsilon$ for some $r \in \mathbb{R}$. Then r is rational and if $r = u/v$ with $u,v \in \mathbb{N}$ and $\operatorname{hcf}(u,v) = 1$, $v \leq a(n+1)/\varepsilon$.*

Proof. Assume that r is irrational. There are infinitely many pairs $(p,q) \in \mathbb{N}^2$ such that $0 < |aq - rp| < \varepsilon/(n+2)$ and in particular, there is a large integral zero $(p_1,q_1) \in \mathbb{N}^2$ of P in that range. Since all pairs (kp_1,kq_1) for $0 \leq k \leq n+1$ are also zeroes of P, $q_1x - p_1y$ divides P. Repeating the process for $n+1$ distinct integral solutions $\{(p_i,q_i)\}_{1\leq i\leq n+1}$ of P in the range $0 < |aq - rp| < \varepsilon/(n+2)$ yields a contradiction.

Now write $r = u/v$ with $\operatorname{hcf}(u,v) = 1$. Fix $k \in \mathbb{N}$ and let $(p,q) \in \mathbb{N}^2$ be an integral solution of $ay - rx = ak/v$. Since $a(q+lu) - r(p+alv) = ak/v$ for all $l \in \mathbb{N}$, if $ak/v < \varepsilon$, $ay - rx - ak/v$ divides P. The degree of P is bounded by n, hence there are at most n such values of k and $a(n+1)/v \geq \varepsilon$. □

Theorem 3.14 (Rationality theorem)**.** *Let (X,Δ) be a klt pair such that $K_X + \Delta$ is not nef and let $a > 0$ be an integer such that $a(K_X + \Delta)$ is Cartier. Let H be a nef and big Cartier divisor and define:*

$$r = r(H) = \max\{t \in \mathbb{R} : H + t(K_X + \Delta) \text{ is nef}\}.$$

Then there are positive integers u and v such that $r = u/v$ and

$$0 < v \leq a(\dim X + 1).$$

Proof. We follow closely the proof given in [KM98].

Step 1. We may assume that the divisor H is basepoint free. Indeed, Theorem 3.4 shows that for $m \gg 0$ and for some $c,d \in \mathbb{N}$, $H' = m(cH + da(K_X + \Delta))$ is basepoint free. Moreover, $r(H)$ and $r(H')$ satisfy

$$r(H) = \frac{r(H') + mda}{mc}.$$

If the denominator of $r(H')$ is v', the denominator v of $r(H)$ divides mcv'. Since c and m can be chosen arbitrarily large, v divides v' and $0 < v \leq v' < a(\dim X + 1)$.

Step 2. For $(p,q) \in \mathbb{N}^2$, define

$$L(p,q) = \operatorname{Bs}|pH + qa(K_X + \Delta)|.$$

Observe that for p, q sufficiently large and such that $0 < aq - rp < \varepsilon$, the sets $L(p, q)$ stabilise. This follows from the Noetherian condition because if (p', q') is such that $0 < aq' - rp' < \varepsilon$ and $p/q < p'/q'$, then $L(p', q') \subset L(p, q)$. Denote by L_0 their limit and set

$$I = \{(p, q) \in \mathbb{Z}^2 \colon 0 < aq - rp < 1 \text{ and } L(p, q) = L_0\}.$$

Since $|pH + qa(K_X + \Delta)|$ is not nef, and in particular not basepoint free, the set L_0 is not empty.

Step 3. Assume that $r = r(H)$ is not rational. Let $f \colon Y \to X$ be a log resolution of (X, Δ). Consider the divisors:

$$D_1 = f^*H, \quad D_2 = f^*(a(K_X + \Delta)), \quad A = K_Y - f^*(K_X + \Delta).$$

Since $\lceil A \rceil$ is f-exceptional and effective, we have

$$H^0(Y, xD_1 + yD_2 + \lceil A \rceil) \simeq H^0(Y, xH + ya(K_X + \Delta)).$$

Let $(x, y) \in \mathbb{N}^2$ be such that $0 < ay - rx < 1$. Then $xD_1 + yD_2 + A - K_Y$ is nef and big, and by Kawamata-Viehweg,

$$P(x, y) = \chi(xD_1 + yD_2 + \lceil A \rceil) = h^0(xD_1 + yD_2 + \lceil A \rceil)$$

is a polynomial of degree at most $\dim X$ and is non-trivial by Non-Vanishing. By Lemma 3.13, there are arbitrarily large integral points $(p, q) \in \mathbb{N}^2$ such that $0 < aq - rp < 1$ and $P(p, q) \neq 0$. Therefore for all $(p, q) \in I$, $|pH + qa(K_X + \Delta)|$ is not empty and hence $L_0 \neq X$.

Let S be a general section of $|pH + qa(K_X + \Delta)|$. The pair $(X, \Delta + S)$ is not klt; define

$$c = \min\{t \in \mathbb{R} : (X, \Delta + tS) \text{ is not klt}\} \leq 1.$$

Let W a minimal non-klt centre of the pair $(X, \Delta + cS)$. By Lemma 2.16, we may assume that W is exceptional. Nadel vanishing shows that the map

$$H^0(Y, p'H + q'a(K_X + \Delta)) \to H^0(W, (p'H + q'a(K_X + \Delta))_{|W})$$

is surjective, where the pair (p', q') is chosen so that $aq' - rp' < aq - rp$. This is a contradiction as in the proof of Theorem 3.4. This proves that r is rational.

Step 4. Let $r = u/v$, where $u, v \in \mathbb{N}$ and $\operatorname{hcf}(u, v) = 1$. Assume that $v > a(\dim X + 1)$.

Lemma 3.13 with $\varepsilon = 1$ shows that there are arbitrarily large positive integers p, q with $0 < aq - rp < 1$ such that $P(p, q) \neq 0$. The linear system $|pH + qa(K_X + \Delta)|$ is not empty for all $(p, q) \in I$. We proceed as in Step 3. $\square$

4. Log terminal and log canonical models

In this section, we define minimal and canonical models of varieties. For pairs, we introduce *log canonical* and *log terminal* models.

Definition 4.1. Let $f\colon (X,\Delta)\to Z$ be a projective morphism.

1. The pair (X,Δ) is a *log terminal* (lt) model over Z if it has dlt singularities and if $K_X+\Delta$ is f-nef.
2. The pair (X,Δ) is a *log canonical* (lc) model over Z if it has lc singularities and if $K_X+\Delta$ is f-ample.

Definition 4.2. A *log terminal model* of a dlt pair $f\colon (X,\Delta_X)\to Z$ is a commutative diagram

$$\begin{tikzcd}
(X,\Delta_X) \arrow[rr, dashed, "\varphi"] \arrow[dr, "f"'] & & (Y,\Delta_Y) \arrow[dl, "g"] \\
& Z &
\end{tikzcd}$$

where φ is birational and:

1. g is projective,
2. φ^{-1} has no exceptional divisors, i.e. φ is contracting,
3. $\Delta_Y=\varphi_*\Delta_X$,
4. $K_Y+\Delta_Y$ is g-nef,
5. $a(E,X,\Delta_X)<a(E,Y,\Delta_Y)$ for every φ-exceptional divisor E.

Remark 4.3.

(a) This definition corresponds to that of *weak canonical* models in [KM98, Definition 3.50]. Note however that [KM98] only assumes that g is proper.
(b) Log terminal models can be defined without requiring φ to be contracting. We add this condition for clarity of the exposition.

Definition 4.4. A *log canonical model* is a diagram as in Definition 4.2, satisfying conditions $(1-3)$ above and:

4′. $K_Y+\Delta_Y$ is g-ample,
5′. $a(E,X,\Delta_X)\le a(E,Y,\Delta_Y)$ for every φ-exceptional divisor E.

A *weak log canonical model* is a diagram as in Definition 4.2, satisfying conditions $(1-4)$ and $(5')$.

Lemma 4.5. *Assume conditions* $(1-4)$ *of Definition 4.2. If* $K_X+\Delta_X$ *is* f*-negative, then condition* (5) *automatically holds.*

Proof. Let W be a common resolution of (X, Δ_X) and (Y, Δ_Y).

$$\begin{tikzcd}
& W \arrow[dl, "p"'] \arrow[dr, "q"] & \\
(X, \Delta_X) \arrow[rr, dashed, "\varphi"] & & (Y, \Delta_Y)
\end{tikzcd}$$

Write

$$\begin{aligned} K_W &= p^*(K_X + \Delta_X) + \sum a(E, X, \Delta_X)E \\ &= q^*(K_Y + \Delta_Y) + \sum a(E, Y, \Delta_Y)E, \end{aligned}$$

where E runs through divisors on W. The result follows from the Negativity Lemma since $q^*(K_Y + \Delta_Y) - p^*(K_X + \Delta_X)$ is $(f \circ p)$-nef and non-trivial on φ-exceptional divisors. □

Definition 4.6. Let (X, Δ) be an lc pair and let $f \colon X \to Z$ be a projective morphism. The *canonical ring* of (X, Δ) over Z is

$$R(X, K_X + \Delta) = \bigoplus_{n \geq 0} f_* \mathcal{O}_X(\lfloor n(K_X + \Delta) \rfloor).$$

Lemma 4.7. *Let (X, Δ_X) be a dlt (respectively klt, lc) pair over Z.*

1. *A log terminal (respectively log canonical) model (Y, Δ_Y) of (X, Δ_X) is dlt (respectively klt, lc).*
2. *If (Y, Δ_Y) is a log terminal or log canonical model of (X, Δ_X), then $R(Y, K_Y + \Delta_Y) = R(X, K_X + \Delta_X)$.*
3. *If (Y, Δ_Y) is a log canonical model of (X, Δ_X), then*
$$Y \simeq \operatorname{Proj}_Z R(X, K_X + \Delta_X).$$
In particular, there is a unique log canonical model of (X, Δ_X).
4. *Any two log terminal models of (X, Δ_X) are isomorphic in codimension* 1.

Proof. These are consequences of the Negativity Lemma. The main difficulty is to show that discrepancies increase in flips or divisorial contractions of dlt pairs [KM98, Corollary 3.44]. □

The following definition and lemma resolve the issue raised in Remark 3.8.

Definition 4.8. Let $\varphi\colon (X, \Delta_X) \to Z$ be a flipping contraction, i.e.:

1. φ is a small projective morphism,
2. $-(K_X + \Delta_X)$ is φ-ample,
3. $\rho(X/Z) = 1$.

A *flip of* φ is a commutative diagram

$$\begin{tikzcd}
(X, \Delta_X) \arrow[rr, dashed] \arrow[dr, "\varphi"'] & & (X^+, \Delta_{X^+}) \arrow[dl, "\varphi^+"] \\
& Z &
\end{tikzcd}$$

where:

1$^+$. φ^+ is small and projective,
2$^+$. $K_{X^+} + \Delta_{X^+}$ is φ^+-ample, with $\Delta_{X^+} = ((\varphi^+)^{-1} \circ \varphi)_* \Delta_X$,
3$^+$. $\rho(X^+/Z) = 1$.

Lemma 4.9. *Let* (X, Δ_X) *be a* $\mathbb{Q}$*-factorial dlt pair and let* $\varphi\colon X \dashrightarrow X'$ *be a divisorial contraction, a Mori fibration or a flip. Then* X' *is* $\mathbb{Q}$*-factorial.*

Proof. [KM98, Propositions 3.36-3.37]. □

Conjecture 4.10 (Existence of flips)**.** *Dlt flips exist.*

Remark 4.11. Let $\varphi\colon (X, \Delta) \to Z$ a flipping contraction. The flip of φ exists if and only if the canonical algebra of (X, Δ) is finitely generated [KM98, Corollary 6.4]. Therefore the problem of existence of flips is local on the base Z, and we often assume that Z is affine.

Conjecture 4.12 (Termination of flips)**.** *There does not exist an infinite sequence of dlt flips.*

Remark 4.13. A $\mathbb{Q}$-factorial dlt pair (X, Δ) is a limit of klt pairs. For all $0 < \delta \ll 1$ the pair $(X, (1-\delta)\Delta)$ is klt as in the proof of [KM98, Proposition 2.43]. The existence of flips of klt contractions therefore implies that of flips of $\mathbb{Q}$-factorial dlt contractions.

Definition 4.14. Let (X, Δ) be a pair.

1. (X, Δ) is a *minimal model* if $K_X + \Delta$ is nef.
2. If (X, Δ) is not nef and there is a Mori fibration $(X, \Delta) \to Y$, then (X, Δ) is a *Mori fibre space*.

Minimal Model Program (MMP). Let (X, Δ_X) be a dlt pair over Z. If existence and termination of flips hold, there is a sequence of birational maps

$$X = X_1 \dashrightarrow X_2 \dashrightarrow \cdots \dashrightarrow X_n = X'$$

over Z, where $(X', \Delta_{X'})$ is either a minimal model or a Mori fibre space. The sequence is obtained as in Flowchart 1.


START
(X,Δ) dlt pair
$K_X+\Delta$ nef?
YES
(X,Δ) minimal model
END
NO
$\phi\colon X\to Z$ contraction of extremal ray
$\dim Z<\dim X$?
YES
$\phi\colon X\to Z$ Mori fibre space
END
NO
codim Exc $\phi\geq 2$?
YES
$\phi^+\colon X^+\to Z$ flip of ϕ
$(X,\Delta):=(X^+,\Delta^+)$
NO
$(X,\Delta):=(Z,\phi_*\Delta)$

FIGURE 1. Flowchart of the Minimal Model Program

Remark 4.15. The MMP can be run for non $\mathbb{Q}$-factorial log canonical pairs (X,Δ_X), where Δ_X is an $\mathbb{R}$-divisor. The MMP for non $\mathbb{Q}$-factorial klt pairs is discussed in [Fuj07].

Proposition 4.16. *Let $f\colon (X,\Delta_X)\to Z$ be a flipping contraction, where (X,Δ_X) is a $\mathbb{Q}$-factorial dlt pair. Assume that $g\colon (Y,\Delta_Y)\to Z$ is a log terminal model of f. Then:*

1. *(Y,Δ_Y) is also a log canonical model of f.*
2. *(Y,Δ_Y) is the flip of f.*

Proof. The morphism g is small because φ^{-1} has no exceptional divisors. We prove that $K_Y+\Delta_Y$ is g-ample.

Let A be a g-ample divisor on Y and denote $A_X=\varphi_*^{-1}A$. Since X is $\mathbb{Q}$-factorial and f is flipping, A_X is $\mathbb{Q}$-Cartier and

$$\lambda(K_X+\Delta_X)+A_X\sim_{\mathbb{Q}} f^*M,$$

where M is a $\mathbb{Q}$-Cartier divisor on Z and $\lambda\in\mathbb{Q}$. Therefore

$$\lambda(K_Y+\Delta_Y)+A\sim_{\mathbb{Q}} g^*M,$$

and λ is non-zero because A is g-ample. The divisor $K_Y + \Delta_Y$ is g-nef if and only if it is g-ample. ☐

MMP with scaling.

Lemma 4.17 ([Bir07, Sho06]). *Let $(X, \Delta + A)$ be a $\mathbb{Q}$-factorial dlt pair over Z such that $K_X + \Delta + A$ is nef. If $K_X + \Delta$ is not nef, there is a $(K_X + \Delta)$-negative extremal ray R and a real number $0 < \lambda \leq 1$ such that $K_X + \Delta + \lambda A$ is nef and trivial on R.*

Proof. Define

$$\lambda = \inf\{-(K_X + \Delta) \cdot \Sigma / A \cdot \Sigma\}$$

as Σ ranges over curves generating extremal rays of (X, Δ); it is enough to show that λ is a minimum. See [Bir07, Sho06] for a proof. ☐

Definition 4.18. Let $(X, \Delta + A)$ be a $\mathbb{Q}$-factorial dlt pair over Z such that $K_X + \Delta + A$ is nef. A $(K_X + \Delta)$-*MMP with scaling of A* is a sequence:

$$(X_1, \lambda_1) \overset{\varphi_1}{\dashrightarrow} \cdots \overset{\varphi_{i-1}}{\dashrightarrow} (X_i, \lambda_i) \overset{\varphi_i}{\dashrightarrow} \cdots ,$$

where for each index i, A_i and Δ_i are the strict transforms of A and Δ on X_i, and

$$\lambda_i = \min\{t \in \mathbb{R} : K_{X_i} + \Delta_i + tA_i \text{ is nef}\}.$$

Each φ_i is a divisorial contraction or a flip associated to a $(K_{X_i} + \Delta_i)$-negative extremal ray $R_i \subset \overline{\mathrm{NE}}(X_i/Z)$ which is $(K_{X_i} + \Delta_i + \lambda_i A_i)$-trivial.

The proof of the existence of minimal models for varieties of log general type given in [BCHM06] rests on several theorems; we recall two of them here.

Theorem 4.19 (Existence of models, [BCHM06, Theorem C]). *Let (X, Δ) be a klt pair projective over Z, where Δ is big over Z. If there is an effective divisor D such that $K_X + \Delta \sim_{\mathbb{R},Z} D$, then (X, Δ) has a log terminal model over Z.*

Theorem 4.20 (Finiteness of models, [BCHM06, Theorem E]). *Let (X, Δ_0) be a klt pair projective over Z. Fix a general ample $\mathbb{Q}$-divisor A over Z and a rational finite dimensional affine subspace $V \subset \mathrm{WDiv}(X)$ that contains Δ_0. Consider the rational polytope*

$$\mathcal{L}_A = \{A + B : B \in V, B \geq 0 \text{ and the pair } (X, A + B) \text{ is lc}\}.$$

Then the set of isomorphism classes of weak log canonical models of (X, Δ) over Z, where $\Delta \in \mathcal{L}_A$, is finite.

Now we are ready to state Special termination with scaling.

Theorem 4.21 ([BCHM06, Lemma 5.1]). *Assume that existence and finiteness of models hold in dimension $n-1$. Let $(X, \Delta + C)$ be an n-dimensional $\mathbb{Q}$-factorial dlt pair such that $K_X + \Delta + C$ is nef. Assume that $\Delta = A + B$, where A is ample, $B \geq 0$ and $\lfloor \Delta \rfloor \subset \operatorname{Supp} B$. Let*

$$(X, \Delta) = (X_1, \Delta_1) \overset{\varphi_1}{\dashrightarrow} (X_2, \Delta_2) \overset{\varphi_2}{\dashrightarrow} \cdots$$

be a sequence of flips for the $(K_X + \Delta)$-MMP with scaling of C. Denote by R_k the exceptional set of the corresponding flipping contraction of φ_k. For k sufficiently large, $R_k \cap \lfloor \Delta_k \rfloor = \emptyset$.

Definition 4.22. Let (X, Δ) be a $\mathbb{Q}$-factorial dlt pair and $f\colon (X, \Delta) \to Z$ a flipping contraction. We say f is a *pre limiting (pl) flipping contraction* if there is an f-negative irreducible component $S \subset \lfloor \Delta \rfloor$.

Theorem 4.23. *Assume that existence and finiteness of models hold in dimension $n-1$. If flips of pl contractions exist in dimension n, then flips of klt contractions exist in dimension n.*

Proof. The proof follows the ideas in [Sho92, K$^+$92, Fuj07].

Let $f\colon (X, \Delta) \to Z$ be a klt flipping contraction. We assume that Z is affine. In this proof subscripts denote proper transforms.

Step 1. Let $\pi\colon W \to Z$ be a resolution of Z and let F_i be generators of $N^1(W/Z)$. We may assume that no $\pi(F_i)$ contains an irreducible component of $f(\Delta)$. Write $Z' = \operatorname{Sing} Z \cup \operatorname{Sing} f(\Delta) \cup f(\operatorname{Exc} f)$. Note that $\operatorname{codim}_Z Z' \geq 2$, and let $\mathcal{I}_{Z'}$ be the ideal sheaf of Z'. Let $i\colon Z \to W$ be a compactification of Z with a very ample line bundle $\mathcal{L}$ on W such that $\mathcal{L}_{|Z} = \mathcal{O}_Z$ and $\mathcal{F} = i_* \mathcal{I}_{Z'} \otimes \mathcal{L}$ is globally generated. If H is the restriction of a general global section of $\mathcal{F}$ to Z, then H is reduced and $f^*H = H_X$ since f is small. Note that H_X and Δ have no common components. By construction, if $\rho\colon W' \to Z$ is any $\mathbb{Q}$-factorial model of Z, the group $N^1(W'/Z)$ is generated by the components of $H_{W'}$ and by the ρ-exceptional divisors.

Let $h\colon Y \to X$ be a log resolution of the pair $(X, \Delta + H_X)$ such that $f \circ h$ is an isomorphism over $Z \smallsetminus H$. Note that all the h-exceptional divisors are components of h^*H_X. Denote by E_i the exceptional divisors of h and let $E = \sum E_i$. The pair $(Y, \Delta_Y + H_Y + E)$ is $\mathbb{Q}$-factorial and dlt over Z. Consider a very ample divisor A on Y which is general in $|A|$ such that $K_X + \Delta_Y + H_Y + E + A$ is nef. Therefore replacing H by $H + (f \circ h)_* A$ we may assume that $K_X + \Delta_Y + H_Y + E$ is nef.

Step 2. We now run the $(K_{\overline{Y}} + \Delta_{\overline{Y}} + E_{\overline{Y}})$-MMP with scaling of $H_{\overline{Y}}$. We construct a sequence

$$(\overline{Y}_1, \Delta_{\overline{Y}_1} + E_{\overline{Y}_1}) \overset{\chi_1}{\dashrightarrow} \cdots \overset{\chi_{i-1}}{\dashrightarrow} (\overline{Y}_i, \Delta_{\overline{Y}_i} + E_{\overline{Y}_i}) \overset{\chi_i}{\dashrightarrow} \cdots,$$

where $\overline{Y}_1 = \overline{Y}$, and each χ_i is a divisorial contraction or a pl flip. Denote by $h_i\colon Y_i \to Z$ the induced map.

If $K_{\overline{Y}_i} + \Delta_{\overline{Y}_i} + E_{\overline{Y}_i}$ is not nef, let R_i be an extremal ray as in Lemma 4.17. Let D_i be any curve whose class is in R_i, in particular $H_{\overline{Y}_i} \cdot D_i > 0$. Since D_i is h_i-exceptional, the definition of H and $h_i^*H \cdot D_i = 0$ imply $D_i \cdot E_{\overline{Y}_i} < 0$. The curve D_i therefore intersects a component of $\lfloor E_{\overline{Y}_i} + \Delta_{\overline{Y}_i} \rfloor = E_{\overline{Y}_i}$ negatively. If the contraction associated to R_i is small, it is pl flipping. Define $\chi_i\colon \overline{Y}_i \dashrightarrow \overline{Y}_{i+1}$ to be the contraction of R_i if it is divisorial, or its flip if it is small.

Theorem 4.21 shows that after finitely many steps we obtain a pair $\tilde{g}\colon (\widetilde{Y}, \Delta_{\widetilde{Y}} + E_{\widetilde{Y}}) \to Z$ such that $K_{\widetilde{Y}} + \Delta_{\widetilde{Y}} + E_{\widetilde{Y}}$ is $\tilde{g}$-nef.

Step 3. Consider a common resolution

$$\begin{tikzcd} & W \arrow[dl, "p"'] \arrow[dr, "q"] & \\ (X, \Delta) \arrow[rr, dashed] & & (\widetilde{Y}, \Delta_{\widetilde{Y}} + E_{\widetilde{Y}}) \end{tikzcd}$$

We have

$$\begin{aligned} K_W &= p^*(K_X + \Delta) - p_*^{-1}\Delta + E_{p,q} + E_p \\ &= q^*(K_{\widetilde{Y}} + \Delta_{\widetilde{Y}} + E_{\widetilde{Y}}) - q_*^{-1}\Delta_{\widetilde{Y}} - q_*^{-1}E_{\widetilde{Y}} + \widetilde{E}_{p,q} + \widetilde{E}_q, \end{aligned}$$

where the divisors $E_{p,q}$ and $\widetilde{E}_{p,q}$ are exceptional for p and q, E_p is exceptional for p but not for q, and $\widetilde{E}_q$ is exceptional for q but not for p. Observe that $q_*^{-1}E_{\widetilde{Y}}$ is exceptional for p but not for q.

The divisor $q^*(K_{\widetilde{Y}} + \Delta_{\widetilde{Y}} + E_{\widetilde{Y}}) - p^*(K_X + \Delta)$ is $(\tilde{g} \circ q)$-nef, and $p_*^{-1}\Delta - q_*^{-1}\Delta_{\widetilde{Y}}$ is effective and exceptional for q but not for p. The Negativity Lemma over Z shows that

$$E_{p,q} < \widetilde{E}_{p,q} \quad \text{and} \quad E_p + q_*^{-1}E_{\widetilde{Y}} \leq 0. \tag{2}$$

Since (X, Δ) is klt, we have $\lceil E_p \rceil \geq 0$, and as $q_*^{-1}E_{\widetilde{Y}}$ is reduced, this implies $E_{\widetilde{Y}} = 0$. In particular, the map $\widetilde{g}$ is small and (2) implies that $(\widetilde{Y}, \Delta_{\widetilde{Y}})$ is klt. The pair $(\widetilde{Y}, \Delta_{\widetilde{Y}})$ is a log terminal model of (X, Δ), and by Proposition 4.16 this is the required flip. □

Remark 4.24. The existence of pl flips can be reduced to the case where the pair $(X, S + B)$ is plt, where S is irreducible as in [Cor07, Remark 2.2.21].

5. RESTRICTED ALGEBRAS AND ADJOINT ALGEBRAS

The rest of the paper is devoted to the proof of Conjecture 4.10 under certain conditions. More precisely, we prove that klt flips exist in dimension n if existence and finiteness of models hold in dimension $n-1$. As is explained in Remark 4.11, the existence of flips is local on the base; for the rest of the paper we consider varieties that are projective over affine varieties.

We work with graded algebras of rational functions $R = \bigoplus_{n\geq 0} R_n \subset k(X)[T]$, where R_0 is a finitely generated $H^0(X, \mathcal{O}_X)$-algebra.

Definition 5.1. A *truncation* of R is an algebra of the form $R^{(I)} = \bigoplus_{n\geq 0} R_{nI}$ for a positive integer I.

We often use the following simple but important lemma without explicit reference.

Lemma 5.2. *A graded algebra R is finitely generated if and only if its truncation $R^{(I)}$ is finitely generated for any I.*

Proof. [Cor07, Lemma 2.3.3]. □

Definition 5.3. A sequence of effective $\mathbb{R}$-b-divisors $\mathbf{M}_\bullet$ is *superadditive* if $\mathbf{M}_{m+n} \geq \mathbf{M}_m + \mathbf{M}_n$ for all $m, n \geq 0$.

A sequence of effective $\mathbb{R}$-b-divisors $\mathbf{\Delta}_\bullet$ is *concave* if

$$\mathbf{\Delta}_{m+n} \geq \frac{m}{n+m}\mathbf{\Delta}_m + \frac{n}{n+m}\mathbf{\Delta}_n$$

for all $n, m \geq 0$. Convex sequences are defined similarly.

Observe that if $\mathbf{M}_\bullet$ is a superadditive sequence of b-divisors, then there are homomorphisms

$$H^0(X, \mathbf{M}_m) \otimes H^0(X, \mathbf{M}_n) \to H^0(X, \mathbf{M}_{m+n})$$

for all $m, n \geq 0$. This justifies the following definition.

Definition 5.4. A *b-divisorial algebra* on X is the algebra of rational functions

$$R(X, \mathbf{M}_\bullet) = \bigoplus_{m\in\mathbb{N}} H^0(X, \mathbf{M}_m),$$

where $\mathbf{M}_\bullet$ is a superadditive sequence of b-divisors on X.

Lemma 5.5. *Let $\mathbf{M}_\bullet$ be a superadditive sequence of mobile b-divisors on X. The b-divisorial algebra $R = R(X, \mathbf{M}_\bullet)$ is finitely generated if and only if there exists an integer i such that $\mathbf{M}_{ik} = k\mathbf{M}_i$ for all $k \geq 0$.*

Proof. Assume that for some i, $\mathbf{M}_{ik} = k\mathbf{M}_i$ for all $k \geq 0$. Passing to a truncation we may assume that $i = 1$. Let $Y \to X$ be a model such that $\mathbf{M}_1$ descends to Y. Then $R = \bigoplus H^0(Y, i\mathbf{M}_{1Y})$ and the result follows from Zariski's theorem.

Conversely, assume that R is finitely generated. Up to truncation, we may assume that R is generated by $H^0(X, \mathbf{M}_1)$. For each j, take a resolution $Y_j \to X$ such that both $\mathbf{M}_1$ and $\mathbf{M}_j$ descend to Y_j. Since the sequence $\mathbf{M}_\bullet$ is concave we have $H^0(Y_j, j\mathbf{M}_{1Y_j}) \subset H^0(Y_j, \mathbf{M}_{jY_j})$, and the finite generation gives $H^0(Y_j, \mathbf{M}_{jY_j}) = H^0(Y_j, \mathbf{M}_{1Y_j})^j \subset H^0(Y_j, j\mathbf{M}_{1Y_j})$. As $j\mathbf{M}_{1Y_j}$ and $\mathbf{M}_{jY_j}$ are free, $H^0(Y_j, j\mathbf{M}_{1Y_j}) = H^0(Y_j, \mathbf{M}_{jY_j})$ implies $j\mathbf{M}_{1Y_j} = \mathbf{M}_{jY_j}$ and thus $j\mathbf{M}_1 = \mathbf{M}_j$. $\square$

Setup 5.6. Consider a pl flipping contraction

$$\pi\colon (X, \Delta = S + B) \to Z,$$

where S is a prime divisor and $\lfloor B \rfloor = 0$. Recall that $-(K_X + \Delta)$ and $-S$ are π-ample, that X is $\mathbb{Q}$-factorial and $\rho(X/Z) = 1$.

The flip of f exists if and only if the canonical algebra

$$R(X, K_X + \Delta) = \bigoplus_{n \geq 0} H^0(X, \lfloor n(K_X + \Delta) \rfloor)$$

is finitely generated.

Remark 5.7. For a Cartier divisor D and a prime Cartier divisor S, let $\sigma_S \in H^0(X, S)$ be a section such that $\operatorname{div} \sigma_S = S$. From the exact sequence

$$0 \to H^0(X, \mathcal{O}_X(D - S)) \xrightarrow{\cdot \sigma_S} H^0(X, \mathcal{O}_X(D)) \xrightarrow{\rho_{D,S}} H^0(S, \mathcal{O}_S(D))$$

we denote $\operatorname{res}_S H^0(X, \mathcal{O}_X(D)) = \operatorname{Im}(\rho_{D,S})$.

Definition 5.8. The *restricted algebra* of $R(X, K_X + \Delta)$ is

$$R_S(X, K_X + \Delta) = \bigoplus_{n \geq 0} \operatorname{res}_S H^0(X, \lfloor n(K_X + \Delta) \rfloor).$$

Lemma 5.9. *Under assumptions of Setup 5.6, $R(X, K_X + \Delta)$ is finitely generated if and only if $R_S(X, K_X + \Delta)$ is finitely generated.*

Proof. We will concentrate on sufficiency, since necessity is obvious.

By [KMM87, Lemma 3-2-5], numerical and linear equivalence over Z coincide. Since $\rho(X/Z) = 1$, and both S and $K_X + \Delta$ are π-negative, there exists a positive rational number r such that $S \sim_{\mathbb{Q},\pi} r(K_X + \Delta)$. By considering open subvarieties of Z we can assume that $S - r(K_X + \Delta)$ is $\mathbb{Q}$-linearly equivalent to a pullback of a principal divisor.

Let φ be a rational function with a zero of order one along S and set $D = S - \operatorname{div} \varphi$; in particular $S \not\subset \operatorname{Supp} D$. Therefore

$$D \sim_{\mathbb{Q}} r(K_X + \Delta),$$

and the rings $R(X, D)$ and $R(X, K_X + \Delta)$ share a common truncation. As $\varphi \in H^0(X, D)$, it is enough to show that φ generates the kernel

$$K = \bigoplus_{n \geq 0} \ker \big(H^0(X, nD) \to H^0(S, nD_{|S})\big).$$

If $\varphi_n \in H^0(X, nD) \cap K$, then $\operatorname{div} \varphi_n + nD - S \geq 0$. Writing $\varphi_n = \varphi\varphi'_n$ for some $\varphi'_n \in k(X)$, we have

$$\operatorname{div} \varphi'_n + (n-1)D = \operatorname{div} \varphi_n - \operatorname{div} \varphi + (n-1)D \geq 0,$$

and therefore $\varphi'_n \in H^0(X, (n-1)D)$. $\square$

Definition 5.10. Let X be a variety that is projective over an affine variety Z. Let $\mathbf{\Delta}_m$ be an effective concave sequence of $\mathbb{Q}$-b-divisors on X such that each $K_X + \mathbf{\Delta}_{mX}$ is $\mathbb{Q}$-Cartier, and assume that there is a positive integer I such that $mI\mathbf{\Delta}_m$ is integral for every m.

An *adjoint algebra* is a b-divisorial algebra

$$R(X, \mathbf{N}_\bullet) = \bigoplus_{m \geq 0} H^0(X, \mathbf{N}_m)$$

where $\mathbf{N}_m = mI(\mathbf{K}_X + \mathbf{\Delta}_m)$, such that:

- $\mathbf{N}_m$ is good on X for every $m \geq 0$, that is $\mathbf{N}_m \geq \overline{\mathbf{N}_{mX}}$,
- $\lim\limits_{m \to \infty} \mathbf{\Delta}_m = \sup\limits_{m \geq 0} \mathbf{\Delta}_m = \mathbf{\Delta}$ exists as a real b-divisor on X.

Remark 5.11. Any truncation of an adjoint algebra is an adjoint algebra.

Lifting Lemma 5.12 (Hacon-M$^\mathrm{c}$Kernan). *Let $(Y, T + B)$ be a log smooth plt pair that is projective over an affine variety Z, where T is irreducible and B is a $\mathbb{Q}$-divisor with $\lfloor B \rfloor = 0$. Assume that*

(1) $B \sim_{\mathbb{Q}} A + C$, where A is ample, $C \geq 0$ and $T \not\subset \operatorname{Supp} C$,
(2) there are a positive integer p and a divisor $N \in |p(K_Y + T + B)|$ such that N meets properly all the intersections of the components of $T + B$.

Then for every m such that $m(K_Y + T + B)$ is integral, the map

$$H^0(Y, m(K_Y + T + B)) \to H^0(T, m(K_T + B_{|T}))$$

is surjective.

Proof. See [HM07]. $\square$

Remark 5.13. The assumptions of the Lifting Lemma are birational. The first assumption is related to B being big; in our setting it is not restrictive. The second assumption, however, is difficult to arrange. Example 5.14 shows that the statement cannot be expected to hold without both assumptions being satisfied.

Example 5.14. Let Y be a surface, let T be a nonsingular curve on Y and let $E \subset Y$ be a (-1)-curve meeting T at exactly one point P. Denote $B = dE$, where $0 < d < 1$. Let $f\colon Y \to X$ be the contraction of E and let $S = f(T)$. Then

$$K_Y + T + B = f^*(K_X + S) + dE, \tag{3}$$

and

$$\begin{tikzcd}
H^0(Y, m(K_Y + T + B)) \arrow[r, "r"] \arrow[d, "\simeq"'] & H^0(T, m(K_T + dP)) \\
H^0(X, m(K_X + S)) \arrow[r] & H^0(T, mK_T)) \arrow[u, hook, "\neq"']
\end{tikzcd}$$

In general, r is not surjective. By (3), $E \subset \operatorname{Bs}|m(K_Y + T + B)|$ and the assumption (2) of the Lifting Lemma is not satisfied.

Definition-Lemma 5.15. Let (X, Δ) be a log pair. There is a b-divisor $\mathbf{B} = \mathbf{B}(X, \Delta)$ on X such that for all models $f\colon Y \to X$, we can write uniquely

$$K_Y + B_Y = f^*(K_X + \Delta) + E_Y,$$

where B_Y and E_Y are effective with no common components and E_Y is f-exceptional. We call $\mathbf{B}$ the *boundary* b-divisor of the pair (X, Δ).

Proof. Let $g\colon Y' \to X$ be a model such that there is a proper birational morphism $h\colon Y' \to Y$. Pushing forward $K_{Y'} + B_{Y'} = g^*(K_X + \Delta) + E_{Y'}$ via h_* yields

$$K_Y + h_*B_{Y'} = f^*(K_X + \Delta) + h_*E_{Y'},$$

and since $h_*B_{Y'}$ and $h_*E_{Y'}$ have no common components, $h_*B_{Y'} = B_Y$. □

Lemma 5.16. *Let (X, Δ) be a log canonical pair. There exists a log resolution $Y \to X$ such that the components of $\{\mathbf{B}_Y\}$ are disjoint.*

Proof. See [KM98, Proposition 2.36] or [HM05, Lemma 6.7]. □

Corollary 5.17. *If (X, Δ) is a klt pair, there are only finitely many valuations E with $a(E, K_X + \Delta) < 0$.*

Proof. Follows from Lemma 5.16 and [KM98, Proposition 2.31]. □

For any two divisors $P = \sum p_i E_i$ and $Q = \sum q_i E_i$ set

$$P \wedge Q = \sum \min\{p_i, q_i\} E_i.$$

Definition 5.18. Let (X,Δ) be a pair such that $I(K_X+\Delta)$ is Cartier and the linear system $|I(K_X+\Delta)|$ is not empty. For a log resolution $f\colon Y\to X$, let

$$mI(K_Y+\mathbf{B}_Y)=M_{mY}+F_{mY}$$

be the decomposition into mobile and fixed parts and set

$$B_{mY}=\mathbf{B}_Y-\mathbf{B}_Y\wedge(F_{mY}/mI).$$

Then

$$\operatorname{Mob}(mI(K_Y+B_{mY}))=M_{mY}=\operatorname{Mob}(mIf^*(K_X+\Delta)).$$

Theorem 5.19. *Let (X,Δ) be a log canonical pair such that $I(K_X+\Delta)$ is Cartier and the linear system $|I(K_X+\Delta)|$ is not empty. Fix a log resolution $f\colon Y\to X$ such that the support of*

$$\operatorname{Fix}|I(K_Y+\mathbf{B}_Y)|+\operatorname{Exc}f+\mathbf{B}_Y$$

is simple normal crossings. There is an effective sequence $\mathbf{B}_\bullet$ of b-divisors on Y such that $\mathbf{B}_m\le\mathbf{B}$ is the smallest b-divisor such that for all models $f\colon Z\to Y$ we have

$$H^0(Z,mI(K_Z+\mathbf{B}_{mZ}))=H^0(Z,mI(K_Z+\mathbf{B}_Z)).$$

The b-divisor $\mathbf{K}_Y+\mathbf{B}_m$ is good on any log resolution $Y_m\to Y$ where M_{mY_m} is free, and the sequence $\mathbf{B}_\bullet$ is concave after truncation.

Proof. Step 1. Fix m and set $\mathbf{B}'=\mathbf{B}(Y,\mathbf{B}_Y)$. Let $g\colon Z\to Y$ be a log resolution and denote $h=g\circ f$. Let

$$K_Z+\mathbf{B}'_Z=g^*(K_Y+\mathbf{B}_Y)+E'_Z,$$

and define M'_{mZ},F'_{mZ} and B'_{mZ} as in Definition 5.18.

Firstly, since $K_Z+\mathbf{B}'_Z=h^*(K_X+\Delta)+g^*E_Y+E'_Z$, we have $\mathbf{B}'_Z=\mathbf{B}_Z+E$ where E is effective. Furthermore,

$$\begin{aligned}M'_{mZ}&=\operatorname{Mob}(mIg^*(K_Y+\mathbf{B}_Y))\\&=\operatorname{Mob}(mIh^*(K_X+\Delta)+mIg^*E_Y)=M_{mZ},\end{aligned}$$

so that

$$mIE=mI(K_Z+\mathbf{B}'_Z)-mI(K_Z+\mathbf{B}_Z)=F'_{mZ}-F_{mZ}.$$

This implies $B_{mZ}=B'_{mZ}$. Now we have

$$mI(K_Y+\mathbf{B}_Y)=g_*(mI(K_Z+\mathbf{B}'_Z))=g_*M'_{mZ}+g_*F'_{mZ},$$

and since $g_*M'_{mZ}=g_*M_{mZ}=M_{mY}$, we obtain $g_*F'_{mZ}=F_{mY}$. Therefore

$$\begin{aligned}g_*B_{mZ}=g_*B'_{mZ}&=g_*\mathbf{B}'_Z-g_*\big(\mathbf{B}'_Z\wedge(F'_{mZ}/mI)\big)\\&=\mathbf{B}_Y-\mathbf{B}_Y\wedge(F_{mY}/mI)=B_{mY},\end{aligned}$$

and $\mathbf{B}_m$ is a well-defined b-divisor.

Step 2. Fix Y_m as in the assumptions of the Theorem, and let Z_m be another log resolution such that M_{mZ_m} is free. We may assume that there is a birational morphism $g_m\colon Z_m \to Y_m$ such that the support of $\operatorname{Fix}|g_m^*(K_{Y_m}+B_{mY_m})|+g_m^*B_{mY_m}+\operatorname{Exc}g_m$ has simple normal crossings. Let $\Gamma_{mZ_m}=\mathbf{B}(Y_m,B_{mY_m})_{Z_m}$; then

$$K_{Z_m}+\Gamma_{mZ_m}=g_m^*(K_{Y_m}+B_{mY_m})+E_{mZ_m}, \tag{4}$$

and $\Gamma_{mZ_m}\leq g_m^*B_{mY_m}$ since Z_m and Y_m are smooth. Furthermore,

$$\begin{aligned}\operatorname{Mob}(mI(K_{Z_m}+\Gamma_{mZ_m}))&=\operatorname{Mob}(g_m^*(mI(K_{Y_m}+B_{mY_m})))\\&=g_m^*M_{mY_m}=M_{mZ_m},\end{aligned}$$

and thus $\operatorname{Fix}|mI(K_{Z_m}+\Gamma_{mZ_m})|=g_m^*\operatorname{Fix}|mI(K_{Y_m}+B_{mY_m})|+E_{mZ_m}$.

Step 3. Let F be a common component of $g_m^*B_{mY_m}$ and $g_m^*\operatorname{Fix}|mI(K_{Y_m}+B_{mY_m})|$. We claim that $a(F,Y_m,B_{mY_m})\geq 0$, so that F is a component of E_{mZ_m}.

To this end, note that the centre of F on Y_m is contained in the intersection of components of B_{mY_m} and $\operatorname{Fix}|mI(K_{Y_m}+B_{mY_m})|$. We can compute the discrepancy of F on any model. By [KM98, Lemma 2.45], there is a composite $\rho_n\colon \mathcal{W}_n\to Y_m$ of n blow ups of the centres of F such that F is a divisor on $\mathcal{W}_n$. Let $\sigma\colon \mathcal{W}_n\to\mathcal{W}_{n-1}$ be the last blowup and let $\rho_{n-1}\colon\mathcal{W}_{n-1}\to Y_m$ be the composite of the first $n-1$ blowups; we can assume $c=\operatorname{codim}_{\mathcal{W}_{n-1}}\sigma(F)>1$. Write

$$K_{\mathcal{W}_{n-1}}+B^+_{\mathcal{W}_{n-1}}=\rho_{n-1}^*(K_{Y_m}+B_{mY_m})+B^-_{\mathcal{W}_{n-1}},$$

where $B^+_{\mathcal{W}_{n-1}}$ and $B^-_{\mathcal{W}_{n-1}}$ are effective divisors without common components. Then by [KM98, Lemmas 2.27, 2.30],

$$\begin{aligned}a(F,Y_m,B_{mY_m})&=a(F,W_{n-1},B^+_{W_{n-1}}-B^-_{W_{n-1}})\\&\geq a(F,W_{n-1},B^+_{W_{n-1}}),\end{aligned} \tag{5}$$

and by induction on n, $B^+_{\mathcal{W}_{n-1}}$ and $\operatorname{Fix}|\rho_{n-1}^*(mI(K_{Y_m}+B_{mY_m}))|$ have no common components. But $\sigma(F)\subset\operatorname{Supp}\operatorname{Fix}|\rho_{n-1}^*(mI(K_{Y_m}+B_{mY_m}))|$, so by the simple normal crossings assumption there can be at most $c-1$ components of $B^+_{\mathcal{W}_{n-1}}$ that contain $\sigma(F)$. Denote these components by $\Delta_1,\dots,\Delta_p$, where $p\leq c-1$. Then by [KM98, Lemma 2.29] and since $(\mathcal{W}_{n-1},B^+_{\mathcal{W}_{n-1}})$ is log canonical, we have

$$a(F,\mathcal{W}_{n-1},B^+_{\mathcal{W}_{n-1}})=c-1+\sum_{i=1}^p a(\Delta_i,\mathcal{W}_{n-1},B^+_{\mathcal{W}_{n-1}})\geq 0, \tag{6}$$

and thus (5) and (6) prove the claim.

As a result, Γ_{mZ_m} and $g_m^* \operatorname{Fix}|mI(K_{Y_m} + B_{mY_m})|$ have no common components and thus $\Gamma_{mZ_m} = B_{mZ_m}$. This proves that $\mathbf{K}_Y + \mathbf{B}_m$ is good on Y_m.

Step 4. To prove that $\mathbf{B}_\bullet$ is concave, observe that the sequence of effective divisors F_{mY}/mI is convex. For each component B_i of $\mathbf{B}_Y$, set $k_i = 1$ if $\operatorname{mult}_{B_i} \mathbf{B}_Y \leq \operatorname{mult}_{B_i} F_{mY}/mI$ for all $m \geq 1$; otherwise, since the sequence F_{mY}/mI has a limit, set k_i to be a positive integer such that $\operatorname{mult}_{B_i} \mathbf{B}_Y \geq \operatorname{mult}_{B_i} F_{mY}/mI$ for all $m \geq k_i$. Let $k = \max\{k_i\}$; the sequence $\mathbf{B}_{km,Y}$ is concave. $\square$

Notation 5.20. The sequence $\mathbf{N}_m = mI(\mathbf{K}_Y + \mathbf{B}_m)$, the associated sequence of mobile b-divisors $\mathbf{M}_m$ (see Lemma 2.6) and the *characteristic* sequence $\mathbf{D}_m = \mathbf{M}_m/m$ are defined on Y. We denote by $R(X, \mathbf{N}_\bullet)$, $R(X, \mathbf{M}_\bullet)$ and $R(X, \mathbf{D}_\bullet)$ the same algebra.

The following two theorems establish that a truncation of the restricted algebra is an adjoint algebra.

Theorem 5.21. *Let (X, Δ) be a plt pair such that $I(K_X + \Delta)$ is Cartier, the linear system $|I(K_X + \Delta)|$ is not empty, $S = \lfloor \Delta \rfloor$ is irreducible and is not a component of* $\operatorname{Fix}|I(K_X + \Delta)|$. *Fix a log resolution $f\colon Y \to X$ such that the components of $\{\mathbf{B}_Y\}$ are disjoint and the support of*

$$\operatorname{Fix}|I(K_Y + \mathbf{B}_Y)| + \operatorname{Exc} f + \mathbf{B}_Y$$

is simple normal crossings. Denote $T = \widehat{S}_Y$ and $B = \mathbf{B}_Y - T$.

Then there are b-divisors $\mathbf{B}^0_m$ on T such that $\mathbf{K}_T + \mathbf{B}^0_m$ is good on T, the limit $\mathbf{B}^0 = \lim_{m\to\infty} \mathbf{B}^0_m$ exists as a b-divisor and the sequence $\mathbf{B}^0_\bullet$ is concave after truncation. More precisely, let $Y_m \to Y$ be any log resolution such that M_{mY_m} is basepoint free and set $R_m = \widehat{S}_{Y_m}$. Then $B^0_{mR_m} = (\mathbf{B}_{mY_m} - R_m)_{|R_m}$.

Remark 5.22. We have the following diagram:

$$\begin{tikzcd}
H^0(Y, mI(K_Y + \mathbf{B}_{mY})) \arrow[r, "r"] \arrow[d, "\simeq"'] & H^0(T, mI(K_T + \mathbf{B}^0_{mT})) \\
H^0(X, mI(K_X + \Delta)) \arrow[r] & \operatorname{res}_S H^0(X, mI(K_X + \Delta)) \arrow[u, hook]
\end{tikzcd}$$

Proof. Step 1. We follow the notation set in the proof of Theorem 5.19. Restricting

$$K_{Y_m} + \mathbf{B}'_{Y_m} = \pi_m^*(K_Y + \mathbf{B}_Y) + E'_{Y_m}$$

to R_m gives

$$K_{R_m} + (\mathbf{B}'_{Y_m} - R_m)_{|R_m} = (\pi_{m|R_m})^*(K_T + B_{|T}) + E'_{Y_m|R_m}.$$

Since the components of $B_{|T}$ do not intersect, the pair $(T, B_{|T})$ is terminal by [KM98, Proposition 2.31], thus $(\mathbf{B}'_{Y_m} - R_m)_{|R_m} = (\pi_{m|R_m})_*^{-1}B_{|T}$ and $E'_{Y_m|R_m}$ is exceptional. Therefore, the components of $B^0_{mR_m}$ do not intersect and $(R_m, B^0_{mR_m})$ is terminal.

Let $Q_m = \widehat{S}_{Z_m}$ and recall from Step 3 of the proof of Theorem 5.19 that $\Gamma_{mZ_m} = B_{mZ_m}$, so that the restriction of (4) to Q_m yields

$$K_{Q_m} + B^0_{mQ_m} = (g_{m|Q_m})^*(K_{R_m} + B^0_{mR_m}) + E_{mZ_m|Q_m},$$

and thus $(g_{m|Q_m})_*^{-1}B^0_{mR_m} = B^0_{mQ_m}$ and $E_{mZ_m|Q_m}$ is exceptional. Set $\mathbf{B}^0_m = \widehat{B^0_{mT}}$, where $B^0_{mT} = (\pi_{m|R_m})_*B^0_{mR_m} \leq B_{|T}$. The pair (T, B^0_{mT}) is terminal, and $\mathbf{K}_T + \mathbf{B}^0_m$ is good on T.

Step 2. To prove concavity of $\mathbf{B}^0_\bullet$ observe that the normal crossings assumption on Y_m ensures that

$$B^0_{mT} = B_{|T} - B_{|T} \wedge \big((\pi_{m|R_m})_*F_{mY_m|R_m}/mI\big).$$

Fix positive integers i and j, and assume that $W = Y_i = Y_j = Y_{i+j}$. Denote $V = \widehat{S}_W$ and $\mu \colon V \to T$. Then $F_{iW} + F_{jW} \geq F_{i+j,W}$ implies

$$\mu_*F_{iW|V} + \mu_*F_{jW|V} \geq \mu_*F_{i+j,W|V},$$

and the sequence $(\pi_{m|R_m})_*F_{mY_m|R_m}/mI$ is convex. The sequence $B^0_{km,T}$ is concave for some k as in Step 4 of the proof of Theorem 5.19. Therefore the limit $\mathbf{B}^0$ exists since $\mathbf{B}^0_{mT} \leq B_{|T}$. □

Notation 5.23. On T, we consider the sequences $\mathbf{N}^0_m = mI(\mathbf{K}_T + \mathbf{B}^0_m)$, the associated sequence of mobile b-divisors $\mathbf{M}^0_m$ as in Lemma 2.6 and the *characteristic* sequence $\mathbf{D}^0_m = \mathbf{M}^0_m/m$.

Theorem 5.24. *Under assumptions of Setup 5.6, a truncation of the restricted algebra $R_S(X, K_X + \Delta)$ is an adjoint algebra. More precisely, there is a log resolution $f \colon Y \to X$ such that, if $T = \widehat{S}_Y$, there is a natural sequence of b-divisors $\mathbf{B}^0_\bullet$ on T with*

$$R_S(X, K_X + \Delta)^{(s)} \simeq \bigoplus_{n \geq 0} H^0(T, nsI(K_T + \mathbf{B}^0_{ns,T}))$$

for some positive integers s and I.

Proof. Every divisor on X is big because π is birational and by Kodaira's lemma $B \sim_{\mathbb{Q}} A + C$, where A is ample and $C \geq 0$. Since π is small, every divisor on X is mobile and we can assume that S is not a component of C. For $0 < \varepsilon < 1$ we have

$$B \sim_{\mathbb{Q}} (1-\varepsilon)B + \varepsilon C + \varepsilon A,$$

and $(X, (1-\varepsilon)B + \varepsilon C + \varepsilon A)$ is klt for $\varepsilon \ll 1$. Replacing A by εA and C by $(1-\varepsilon)B + \varepsilon C$ we may assume that $B = A + C$ with A ample and C effective, and such that $S \not\subset \operatorname{Supp} C$. Furthermore, if k is an integer such that kA is very ample, we may assume kA is a very general member of the linear system $|kA|$ so that A is transverse to all other relevant divisors.

Let $f\colon Y \to X$ be a resolution as in Theorem 5.21. Observe that $R^{(I)} = R(Y, \mathbf{N}_\bullet)$, and we will show that $R^{(I)}_S = R(T, \mathbf{N}^0_\bullet)$.

Fix a positive integer m. It will be enough to apply the Lifting Lemma to conclude that the restriction map $H^0(Y, \mathbf{N}_m) \to H^0(T, \mathbf{N}^0_m)$ is surjective.

For condition (1) of the Lifting Lemma, since kA is very general in $|kA|$, $f^*A = f^{-1}_*A$ is free and hence $\mathbf{B}_{mY} - T \geq f^*A$. We can choose a small f-exceptional effective divisor E on Y such that $A' = f^*A - E$ is ample, and $\mathbf{B}_{mY} - T = A' + C'$, where the support of $C' = \mathbf{B}_{mY} - T - f^*A + E$ does not contain T.

We prove that condition (2) of the Lifting Lemma is satisfied. We can assume that $\operatorname{Mob} \mathbf{N}_{mY}$ is free. Since $\operatorname{Fix} |\mathbf{N}_{mY}|$ and $\mathbf{B}_{mY}$ have simple normal crossings supports and have no common components, $\operatorname{Bs} |\mathbf{N}_{mY}|$ does not contain any intersection of components of $\mathbf{B}_{mY}$. This finishes the proof. □

Corollary 5.25. *$\mathbf{M}^0_i = \mathbf{M}_{i|T}$ for every i.*

Proof. Fix i. Let $Y' \to Y$ be a resolution such that, if $W = \widehat{T}_{Y'}$, $\mathbf{M}_i$ descends to Y' and $\mathbf{M}^0_i$ descends to W. Restricting $\mathbf{M}_{iY'} \leq \mathbf{N}_{iY'}$ to W and taking mobile parts shows that $\mathbf{M}_{iY'|W} \leq \mathbf{M}^0_{iW}$, and hence

$$H^0(W, \mathbf{M}_{iY'|W}) \subset H^0(W, \mathbf{M}^0_{iW}).$$

As

$$H^0(W, \mathbf{M}^0_{iW}) = \operatorname{Image}\bigl(H^0(Y', \mathbf{M}_{iY'}) \to H^0(W, \mathbf{M}_{iY'|W})\bigr),$$

$\mathbf{M}_{iY'|W} = \mathbf{M}^0_{iW}$ because both divisors are free. □

Theorem 5.26. *Assume existence and finiteness of models. Let (X, Δ) be a $\mathbb{Q}$-factorial klt pair projective over Z with $K_X + \Delta$ big, and let $V \subset \operatorname{Div}(X)_{\mathbb{R}}$ be a finite-dimensional vector space that contains Δ. Let $\Delta_\bullet$ be a concave sequence with $\lim_{m\to\infty} \Delta_m = \Delta$ and assume there is a positive integer I such that $I(K_X + \Delta_m)$ is integral for every m.*

Then there exist a smooth model $f\colon Y \to X$ *and a positive integer* s *such that*

$$\lim_{m\to\infty} \operatorname{Mob}(msIf^*(K_X+\Delta_m))/m$$

exists and is a semiample divisor.

Proof. By Kodaira's Lemma, $K_X+\Delta \sim_{\mathbb{R},f} A+C$, where A is ample and C is effective. If $\Delta' = \varepsilon A + \varepsilon C + \Delta$ for $0<\varepsilon \ll 1$, the pair (X,Δ') is klt and $K_X+\Delta' \sim_{\mathbb{R},f} (1+\varepsilon)(K_X+\Delta)$. Therefore replacing Δ by Δ' we may assume that Δ is big.

Let $\varphi_i\colon X \dashrightarrow W_i$ be finitely many weak log canonical models corresponding to the polytope around Δ. Let $f\colon Y\to X$ be a common log resolution of the pair (X,Δ) and of the pairs $(W_i,(\varphi_i)_*\Delta)$ for all i, and denote $\pi_i\colon Y\to W_i$ the induced maps. For every $m\gg 0$ there is an index i_m such that $K_{W_{i_m}}+(\varphi_{i_m})_*\Delta_m$ is semiample by Theorem 3.4.

Since W_i have rational singularities, by the proof of [Kaw88, Lemma 1.1], the groups $\operatorname{WDiv}(W_i)/\operatorname{Div}(W_i)$ are finitely generated, and hence finite because each W_i is $\mathbb{Q}$-factorial. Therefore, for $m\gg 0$ there are positive integers s_{i_m} such that $s_{i_m}I(K_{W_{i_m}}+(\varphi_{i_m})_*\Delta_m)$ are Cartier. By Effective Basepoint Freeness, there is a positive integer c such that all $cs_{i_m}I(K_{W_{i_m}}+(\varphi_{i_m})_*\Delta_m)$ are free; set $s = c\prod s_i$.

By the Negativity Lemma, for $m\gg 0$ we can write

$$f^*(K_X+\Delta_m) = \pi_{i_m}^*(K_{W_{i_m}}+(\varphi_{i_m})_*\Delta_m)+E_m,$$

where E_m are effective and exceptional. There is an index i such that $i=i_m$ for infinitely many m and therefore the divisor

$$\begin{aligned}\lim_{m\to\infty}\operatorname{Mob}(msIf^*(K_X+\Delta_m))/m &= \lim_{m\to\infty} sI\pi_i^*(K_{W_i}+(\varphi_i)_*\Delta_m)\\ &= sI\pi_i^*(K_{W_i}+(\varphi_i)_*\Delta)\end{aligned}$$

is nef and thus semiample by Theorem 3.4. □

Definition 5.27. An adjoint algebra $R(X,\mathbf{D}_\bullet)$ is *semiample* if $\mathbf{D} = \lim_{i\to\infty}\mathbf{D}_i$ exists as a real mobile b-divisor.

Theorem 5.28. *Let* $(X,\Delta)\to Z$ *be a pl flipping contraction as in Setup 5.6, where* $\dim X = n$. *Assume existence and finiteness of models in dimension* $n-1$. *Then there is a positive integer* s *such that* $R_S(X,K_X+\Delta)^{(s)}$ *is a semiample adjoint algebra. Moreover, we can assume that there is a model* T *to which all* $\mathbf{M}_m^0$ *descend.*

Proof. Let a log resolution $f\colon Y\to X$ and the associated sequence $\mathbf{B}_\bullet^0$ be as in Theorem 5.21. Let $\mathbf{B}^0 = \lim\limits_{i\to\infty}\mathbf{B}_i^0 \in \mathbf{Div}(T)_{\mathbb{R}}$. By Corollary 5.26 there is a model $W\to T$ and a positive integer s such that $\mathbf{M}^0_{ms,W}$ is free for every m and $\lim\limits_{m\to\infty}\mathbf{M}^0_{ms,W}/m$ is semiample. We are done. □

6. Shokurov's saturation condition

Definition 6.1. Let (X, Δ) be a log pair, let $\mathbf{A} = \mathbf{A}(X, \Delta)$ and let $\mathbf{F}$ be a b-divisor such that $\lceil \mathbf{F} \rceil \geq 0$.

- An integral mobile b-divisor $\mathbf{M}$ on X is $\mathbf{F}$-*saturated* if there is a model $Y \to X$ such that
$$\operatorname{Mob}\lceil \mathbf{M}_W + \mathbf{F}_W \rceil \leq \mathbf{M}_W$$
on all models $W \to Y$. If (X, Δ) is klt and $\mathbf{F} = \mathbf{A}$ then we say that $\mathbf{M}$ is saturated.
- An adjoint algebra $R = R(X, \mathbf{N}_\bullet)$ is (asymptotically) $\mathbf{F}$-saturated if there is a model $Y \to X$ such that on every log smooth model $(W, \mathbf{F}_W)$ over Y, for all $i \geq j > 0$ we have
$$\operatorname{Mob}\lceil j\mathbf{D}_{iW} + \mathbf{F}_W \rceil \leq j\mathbf{D}_{jW}.$$
If (X, Δ) is klt and $\mathbf{F} = \mathbf{A}$ then we say that R is saturated.

Remark 6.2. Any truncation of a saturated adjoint algebra is saturated.

Example 6.3. If M is a divisor on a klt pair (X, Δ), then $\mathbf{M} = \overline{M}$ is saturated. Indeed, for every model $f\colon Y \to X$ we have $f_*\mathcal{O}_Y(f^*M + E) = \mathcal{O}_X(M)$ for an effective and f-exceptional divisor E.

We prove that saturated semiample adjoint algebras on curves re finitely generated; this models the proof in higher dimensions.

Proposition 6.4. *Let $X = \mathbb{A}^1_{\mathbb{C}}$. A saturated adjoint algebra on X is finitely generated.*

Proof. On a curve b-divisors are just ordinary divisors. Consider an adjoint algebra $R = R(X, N_\bullet)$. Since on an affine curve all linear systems are basepoint free, we have $N_i = M_i = iD_i$ and since $A = -\Delta$, the saturation condition reads
$$\lceil jD_i - \Delta \rceil \leq jD_j.$$
This is a componentwise condition, so we can assume that all divisors are supported at a point $P \in \mathbb{A}^1_{\mathbb{C}}$. Let $\Delta = bP$ with $0 < b < 1$ and $D_i = d_iP$, so that
$$R(X, N_\bullet) = \bigoplus_{i \geq 0} H^0(\mathbb{A}^1_{\mathbb{C}}, \mathcal{O}_{\mathbb{A}^1_{\mathbb{C}}}(id_iP)).$$
The sequence d_i satisfies $d_i \geq 0$ and $d_{i+j} \geq \frac{i}{i+j}d_i + \frac{j}{i+j}d_j$, so it has a limit $d = \sup d_i$. The saturation condition when $i \to \infty$ becomes
$$\lceil jd - b \rceil \leq jd_j$$

for all $j > 0$. Denote $q = \lfloor 1/(1-b) \rfloor$.

Claim 6.5. *$q!d \in \mathbb{N}$ and $d = d_j$ if $jd \in \mathbb{N}$.*

In particular if $d = u/v$ with $u, v \in \mathbb{N}$, then $R^{(v)} = R(\mathbb{A}^1_{\mathbb{C}}, uP)$ and R is finitely generated.

To prove the claim, first assume that $d \notin \mathbb{Q}$; then there is $j \in \mathbb{N}$ such that $\{jd\} > b$ and therefore

$$jd_j \leq jd < \lceil jd - b \rceil \leq jd_j,$$

a contradiction. Assume now that $jd \in \mathbb{Z}$, then

$$jd_j \geq \lceil jd - b \rceil = jd \geq jd_j,$$

and therefore $d = d_j$. Choose the smallest such j and set $l = jd$. Then l and j are coprime, so there is $k < j$ such that $kl \equiv -1 \pmod{j}$. But then if $j > q$, $\{kd\} = \{kl/j\} = (j-1)/j > b$ and this is a contradiction as above. □

Lemma 6.6. *Let $Y \to X$ be a log resolution of a pair (X, Δ). Assume that $\mathbf{N}$ is a real b-divisor on X which descends to Y and such that $\operatorname{Supp} \mathbf{N}_Y$ is simple normal crossings. Then $\lceil \mathbf{N} + \mathbf{A} \rceil$ is good on Y.*

Proof. Let $f \colon Y' \to Y$ be a model and let G be any simple normal crossings divisor on Y. The pair $(Y, \lceil G \rceil - G)$ is klt and in particular the divisor

$$E = \lceil K_{Y'} - f^*(K_Y + \lceil G \rceil - G) \rceil$$

is effective and exceptional. Thus $f^*\lceil G \rceil + E = \lceil K_{Y'/Y} + f^*G \rceil$. Consider the divisor $G = \mathbf{N}_Y + \mathbf{A}_Y$ and observe that $\mathbf{A}_{Y'} = f^*\mathbf{A}_Y + K_{Y'/Y}$; this yields:

$$f^*\lceil \mathbf{N}_Y + \mathbf{A}_Y \rceil + E = \lceil f^*\mathbf{N}_Y + \mathbf{A}_{Y'} \rceil = \lceil \mathbf{N}_{Y'} + \mathbf{A}_{Y'} \rceil.$$

□

Lemma 6.7. *Let S be a subvariety of a projective variety X and let $h \colon T \to S$ be a projective birational map. Then there is a projective birational map $f \colon Y \to X$ from a variety Y and a closed immersion $j \colon T \to Y$ such that $f \circ j = h$.*

Furthermore, assume that h is proper but not necessarily projective. Then there exist a projective birational map $f \colon Y \to X$ and a subvariety $T' \subset Y$ such that $f_{|T'}$ factors through h.

Proof. Let $i \colon S \to X$ be the closed immersion. By [Har77, Chapter II, Theorem 7.17] there is a coherent ideal sheaf $\mathcal{J}$ on S such that T is the blowup of S with respect to $\mathcal{J}$. Let Y be the blowup of X with respect to the ideal sheaf $i_*\mathcal{J}$. The conclusion follows from [Har77, Chapter II, Corollary 7.15] since $i^{-1}(i_*\mathcal{J}) \cdot \mathcal{O}_S = \mathcal{J}$.

The second statement follows from Chow's lemma. □

Theorem 6.8. *Let* $(X,\Delta)\to Z$ *be a pl flipping contraction as in Setup 5.6, where* $\dim X=n$. *Assume the MMP with scaling for* $\mathbb{Q}$*-factorial klt pairs with big boundary in dimension* $n-1$. *There is a positive integer* s *such that* $R_S(X,K_X+\Delta)^{(s)}$ *is saturated.*

Proof. Let s and T be a positive integer and a model as in Theorem 5.28; we assume that T sits on a model $f\colon Y\to X$ by Lemma 6.7. Up to truncation we may assume that $s=1$ and it is enough to show that the saturation condition holds on T. Set $\mathbf{A}^*=\widehat{S}+\mathbf{A}$ and $\mathbf{A}^0=\mathbf{A}(S,B_S)$.

Fix integers $i\geq j>0$. Let $\varphi\colon Y'\to Y$ be a log resolution such that $\mathbf{M}_i$ and $\mathbf{M}_j$ descend to Y' and $\mathbf{M}_i^0$ and $\mathbf{M}_j^0$ descend to $W=\widehat{T}_{Y'}$. Furthermore, by Lemma 6.6 and Lemma 2.6 we can assume that $\operatorname{Mob}\lceil j\mathbf{D}_{iY'}+\mathbf{A}^*_{Y'}\rceil$ is free. Let $g=f\circ\varphi$ and observe that $\lceil\mathbf{A}^*_{Y'}\rceil$ is effective and g-exceptional. Denoting $M=I(K_X+S+B)$, since $\mathbf{D}_{iY'}=(1/i)\operatorname{Mob}g^*(iM)$ we have

$$\begin{aligned}\operatorname{Mob}\lceil j\mathbf{D}_{iY'}+\mathbf{A}^*_{Y'}\rceil&\leq\operatorname{Mob}\lceil (j/i)g^*(iM)+\mathbf{A}^*_{Y'}\rceil\\&=\operatorname{Mob}g^*(jM)=j\mathbf{D}_{jY'},\end{aligned}$$

and therefore

$$(\operatorname{Mob}\lceil j\mathbf{D}_{iY'}+\mathbf{A}^*_{Y'}\rceil)_{|W}\leq j\mathbf{D}^0_{jW},$$

since $\mathbf{D}^0_{iW}=\mathbf{D}_{iY'|W}$ the proof of Corollary 5.25.

By adjunction $\mathbf{A}^*_{Y'|W}=\mathbf{A}^0_W$. Taking global sections in the sequence

$$\begin{aligned}0\to{}&\mathcal{O}_{Y'}(\lceil j\mathbf{D}_{iY'}+\mathbf{A}_{Y'}\rceil)\\&\to\mathcal{O}_{Y'}(\lceil j\mathbf{D}_{iY'}+\mathbf{A}^*_{Y'}\rceil)\to\mathcal{O}_W(\lceil j\mathbf{D}^0_{iW}+\mathbf{A}^0_W\rceil)\to 0\end{aligned}$$

we obtain that the restriction map

$$H^0(Y',\lceil j\mathbf{D}_{iY'}+\mathbf{A}^*_{Y'}\rceil)\to H^0(W,\lceil j\mathbf{D}^0_{iW}+\mathbf{A}^0_W\rceil)$$

is surjective since $H^1(Y',\lceil j\mathbf{D}_{iY'}+\mathbf{A}_{Y'}\rceil)=(0)$ by Kawamata-Viehweg vanishing. Then $(\operatorname{Mob}\lceil j\mathbf{D}_{iY'}+\mathbf{A}^*_{Y'}\rceil)_{|W}=\operatorname{Mob}\lceil j\mathbf{D}^0_{iW}+\mathbf{A}^0_W\rceil$ as in the proof of Corollary 5.25, and therefore

$$\operatorname{Mob}\lceil j\mathbf{D}^0_{iW}+\mathbf{A}^0_W\rceil\leq j\mathbf{D}^0_{jW}.$$

Finally by Lemma 6.6 and Lemma 2.6,

$$\begin{aligned}\operatorname{Mob}\lceil j\mathbf{D}^0_{iT}+\mathbf{A}^0_T\rceil&=(g_{|W})_*\operatorname{Mob}\lceil j\mathbf{D}^0_{iW}+\mathbf{A}^0_W\rceil\\&\leq(g_{|W})_*j\mathbf{D}^0_{jW}=j\mathbf{D}^0_{jT}.\end{aligned}$$

This concludes the proof. □

7. Existence of flips

Approximation Lemma 7.1. *Let Y be a nonsingular variety and let $P_1, \dots, P_l$ be free integral divisors on Y. Let $D \in \sum \mathbb{R}_+ P_k$ and assume that D is not a $\mathbb{Q}$-divisor. Let $\sum G_i$ be a reduced divisor containing* $\operatorname{Supp} P_k$ *for every k. Then for every $0 < \varepsilon \ll 1$ there exists a free integral divisor M and a positive integer j such that:*

1. *$\|jD - M\|_\infty < \varepsilon$, where the sup-norm is taken with respect to the basis $\{G_i\}$ of the vector space $\bigoplus \mathbb{R}G_i \subset \operatorname{Div}(Y)$,*
2. *$jD - M$ is not effective.*

Proof. Set $L_\mathbb{Z} = \bigoplus \mathbb{Z}P_k \simeq \mathbb{Z}^l$ and $L_\mathbb{R} = \bigoplus \mathbb{R}P_k \simeq \mathbb{R}^l$, and assume that D is represented in that basis by $\mathbf{d} = (d_1, \dots, d_l) \in \mathbb{R}^l_+$, where not all d_i are rational.

Let $\pi \colon L_\mathbb{R} \to L_\mathbb{R}/L_\mathbb{Z}$ be the projection map and let $A = \mathbb{N}\pi(\mathbf{d}) \subset L_\mathbb{R}/L_\mathbb{Z}$. The closure $\overline{A}$ is a positive dimensional Lie subgroup; denote by $\overline{A}_0$ the positive dimensional real sub-torus that is the connected component of 0. Then $T = \pi^{-1}(\overline{A}_0) \subset L_\mathbb{R}$ is a positive dimensional vector space and T is not a subset of the cone $L_\mathbb{R} \cap \sum \mathbb{R}_+ G_i$. Therefore for every $\varepsilon > 0$ there is a positive integer j and an element $\mathbf{m} = (m_1, \dots, m_l) \in \mathbb{Z}^l$ such that $\|j\mathbf{d} - \mathbf{m}\|_\infty < \varepsilon$ and $j\mathbf{d} - \mathbf{m} \notin L_\mathbb{R} \cap \sum \mathbb{R}_+ G_i$. If $\varepsilon \ll 1$, then $\mathbf{m} \in \mathbb{N}^l$ and thus $\sum m_i P_i$ is free. □

Finally we can prove:

Theorem 7.2. *Let $R = R(Y, \mathbf{N}_\bullet)$ be an adjoint algebra. If R is saturated and semiample then it is finitely generated.*

Proof. Firstly we prove that $\mathbf{D} = \lim\limits_{i \to \infty} \mathbf{D}_i$ is a $\mathbb{Q}$-b-divisor. Assume otherwise. Passing to a resolution we may assume that $\mathbf{D}_Y$ is semiample. Letting $i \to \infty$ in the saturation condition, we have

$$\operatorname{Mob}\lceil j\mathbf{D}_Y + \mathbf{A}_Y \rceil \le j\mathbf{D}_{jY}$$

for all $j > 0$. Let $P_1, \dots, P_l$ be free integral divisors on Y such that $\mathbf{D}_Y \in \sum \mathbb{R}_+ P_k$, and let $G = \sum G_i$ be a reduced divisor containing $\operatorname{Supp} P_k$ for every k. Fix $\varepsilon > 0$ small enough so that $\lceil \mathbf{A}_Y - \varepsilon G \rceil \ge 0$. By the Approximation Lemma there is a free integral divisor M and a positive integer j such that $\|j\mathbf{D}_Y - M\|_\infty < \varepsilon$ and $j\mathbf{D}_Y - M$ is not effective. Then $j\mathbf{D}_Y - M \ge -\varepsilon G$ and

$$\lceil j\mathbf{D}_Y + \mathbf{A}_Y \rceil = \lceil M + \mathbf{A}_Y + (j\mathbf{D}_Y - M) \rceil \ge M + \lceil \mathbf{A}_Y - \varepsilon G \rceil \ge M,$$

so that

$$j\mathbf{D}_Y \ge j\mathbf{D}_{jY} \ge \operatorname{Mob}\lceil j\mathbf{D}_Y + \mathbf{A}_Y \rceil \ge M.$$

This is a contradiction.

Let now j be a positive integer such that $j\mathbf{D}_Y$ is an integral free divisor. Then

$$j\mathbf{D}_{jY} \leq j\mathbf{D}_Y \leq \operatorname{Mob}(j\mathbf{D}_Y + \lceil \mathbf{A}_Y \rceil) \leq j\mathbf{D}_{jY},$$

and thus $\mathbf{D}_Y = \mathbf{D}_{jY}$ and $\mathbf{D} = \mathbf{D}_j$. □

Corollary 7.3. *Assume existence and finiteness of models in dimension $n-1$. Then klt flips exist in dimension n.*

Proof. By Theorem 4.23 and Remark 4.24, it is enough to prove that the flip of a pl flipping contraction $(X, \Delta) \to Z$ exists, where $S = \lfloor \Delta \rfloor$ is irreducible. Since the problem is local on the base we assume that Z is affine, and therefore it is enough to prove that the canonical algebra $R(X, K_X + \Delta)$ is finitely generated. By Theorems 5.24, 5.28 and 6.8, a truncation of the restricted algebra $R_S(X, K_X + \Delta)$ is a saturated semiample adjoint algebra, and thus is finitely generated by Theorem 7.2. □

REFERENCES

[Amb03] Florin Ambro, *Quasi-log varieties*, Tr. Mat. Inst. Steklova **240** (2003), 220–239.

[BCHM06] Caucher Birkar, Paolo Cascini, Christopher D. Hacon, and James M^{c}Kernan, *Existence of minimal models for varieties of log general type*, arXiv:math.AG/0610203v2.

[Bir07] Caucher Birkar, *On existence of log minimal models*, arXiv:math.AG/0706.1792.

[Cor07] Alessio Corti, *3-fold flips after Shokurov*, Flips for 3-folds and 4-folds (Alessio Corti, ed.), Oxford Lecture Series in Mathematics and its Applications, vol. 35, Oxford University Press, 2007, pp. 18–48.

[Fuj07] Osamu Fujino, *Special termination and reduction to pl flips*, Flips for 3-folds and 4-folds (Alessio Corti, ed.), Oxford Lecture Series in Mathematics and its Applications, vol. 35, Oxford University Press, 2007, pp. 63–75.

[Har77] Robin Hartshorne, *Algebraic geometry*, Springer-Verlag, New York, 1977, Graduate Texts in Mathematics, No. 52.

[HM05] Christopher D. Hacon and James M^{c}Kernan, *On the existence of flips*, arXiv:math.AG/0507597.

[HM07] ———, *Extension theorems and the existence of flips*, Flips for 3-folds and 4-folds (Alessio Corti, ed.), Oxford Lecture Series in Mathematics and its Applications, vol. 35, Oxford University Press, 2007, pp. 76–110.

[HM08] ———, *Existence of minimal models for varieties of log general type II*, arXiv:0808.1929v1.

[Iit82] Shigeru Iitaka, *Algebraic geometry*, Graduate Texts in Mathematics, vol. 76, Springer-Verlag, New York, 1982.

[K$^+$92] János Kollár et al., *Flips and abundance for algebraic threefolds*, Astérisque 211, Soc. Math. France, Paris, 1992.

[Kaw88] Yujiro Kawamata, *Crepant blowing-up of* 3*-dimensional canonical singularities and its application to degenerations of surfaces*, Ann. of Math. (2) **127** (1988), no. 1, 93–163.

[KM98] János Kollár and Shigefumi Mori, *Birational geometry of algebraic varieties*, Cambridge Tracts in Mathematics, vol. 134, Cambridge University Press, Cambridge, 1998.

[KMM87] Yujiro Kawamata, Katsumi Matsuda, and Kenji Matsuki, *Introduction to the minimal model problem*, Algebraic geometry, Sendai, 1985, Adv. Stud. Pure Math., vol. 10, North-Holland, Amsterdam, 1987, pp. 283–360.

[Kol93] János Kollár, *Effective base point freeness*, Math. Ann. **296** (1993), no. 4, 595–605.

[Kol07] ______, *Kodaira's Canonical bundle formula and subadjunction*, Flips for 3-folds and 4-folds (Alessio Corti, ed.), Oxford Lecture Series in Mathematics and its Applications, vol. 35, Oxford University Press, 2007, pp. 134–162.

[Laz04] Robert Lazarsfeld, *Positivity in algebraic geometry. II*, Ergebnisse der Mathematik und ihrer Grenzgebiete, vol. 49, Springer-Verlag, Berlin, 2004.

[Sho92] Vyacheslav V. Shokurov, *Three-dimensional log perestroikas*, Izv. Ross. Akad. Nauk Ser. Mat. **56** (1992), no. 1, 105–203.

[Sho03] ______, *Prelimiting flips*, Tr. Mat. Inst. Steklova **240** (2003), 82–219.

[Sho06] ______, *Letters of a Bi-Rationalist: VII. Ordered termination*, arXiv:0607822.

[Zar62] Oscar Zariski, *The theorem of Riemann-Roch for high multiples of an effective divisor on an algebraic surface*, Ann. of Math. **76** (1962), no. 3, 560–615, with an appendix by David Mumford.

Department of Mathematics, Imperial College London, Huxley Building, 180 Queen's Gate, London SW7 2AZ, United Kingdom
E-mail address: `A.Corti@imperial.ac.uk`

Department of Pure Mathematics and Mathematical Statistics, University of Cambridge, Wilberforce Road, Cambridge CB3 0WB, United Kingdom
E-mail address: `A.S.Kaloghiros@dpmms.cam.ac.uk`

Department of Pure Mathematics and Mathematical Statistics, University of Cambridge, Wilberforce Road, Cambridge CB3 0WB, United Kingdom
E-mail address: `V.Lazic@dpmms.cam.ac.uk`